TO EVALUATE THE EFFECT OF CONTINUOUS IRRIGATION AND SUCTION TECHNIQUE BY USING ANTIBIOTIC SOLUTION IN THE TREATMENT OF CHRONIC OSTEOMYELITIS

THESIS FOR MASTER OF SURGERY (ORTHOPAEDICS)



D1135

BUNDELKHAND UNIVERSITY
JHANSI (U.P.)

DEPARTMENT OF CRTHOPAEDIC SURGE M.L.B. MEDICAL COLLEGE, HOSPIT JHANSI (U.P.).

CERTIFICATE

This is to certify that the work entitled "TO EVALUATE THE EFFECT OF CONTINUOUS IRRIGATION AND SUCTION TECHNIQUE BY USING ANTIBIOTIC SOLUTION IN THE TREATMENT OF CHRONIC OSTEOMYELITIS" has been carried out by Dr. Niraj Jain himself in this department.

He has put in the necessary stay in the department as required by the regulation of Bundelkham University.

(P.K. DABRAL)

Professor and Head, Department of Orthopsedic Surge M.L.D. Medical College, Jhansi (U.P.)

DEPARTMENT OF CRITHOPAEDIC SURGERY. M.L.B. MEDICAL COLLEGE, HOSPITAL, JHANSI (U.P.)

CERTIFICATE

This is to certify that the work entitled *TO EVALUATE THE EFFECT OF CONTINUOUS IRRIGATION AND SUCTION TECHNIQUE BY USING ANTIBIOTIC SOLUTION IN THE TREATMENT OF CHRONIC OSTEOMYELITIS", which is being submitted as a thesis for M.S. (Orthopaedic Surgery) was carried out by Dr. Niraj Jain, under my constant supervision and quidance.

The techniques embodied in this work were undertaken by the candidate himself. The results and observations were checked and verified by me periodical

(R. P. TRIPATHI)

M.S.

R. P. Topath

Reader, Department of Orthopaedics M.L.B. Medical College, Jhansi(U.P.)

(GUIDE)

CERTIFICATE

This is to certify that Dr. Miraj Jain
has worked on "TO EVALUATE THE EFFECT OF CONTINUOUS
IRRIGATION AND SUCTION TECHNIQUE BY USING ANTIBIOTIC
SOLUTION IN THE TREATMENT OF CHRONIC OSTEOMYELITIS"
under my direct supervision and guidance.

His results and observations have been checked and verified by me from time to time.

(A. K. GUPTA)

Department of Radiology.
M.L.B. Medical College.
Jhansi (U.P.)

(CO-GUIDE)

I am sure, I can never manage to express my sincere gratitude towards all who have ment so much in the formation of this project. Yet, I shall try.

This has indeed been a rare privilege to have had an opportunity to work under the inspiring guidence of Dr. R.P. Tripathi, M.S., Reader in Orthopeedics, M.L.B. Medical College, Jhansi. It has all been because of his constructive criticism and persistant indeavour to get the best from his student. To him, I owe an inexpressible gratitude.

I have no words to express my regards to my exteemed teacher Dr. A.K. Gupta, M.D., Lecturer in Radiology, M.L.B. Medical College, Jhansi for his keen interest, constant encouragement, timely suggestions, criticism and unlimited help.

I feel, I owe this work to my learned and experienced teacher Prof. P.K. Dabral, N.S., Head, department of Orthopsedics, H.L.B. Medical College, Jhansi. This work could have never been completed had he not shown the path through the unsurmountable obstacles.

I must express my grateful thanks to Dr. D. K.
Gupta, M.S. and Dr. Amurag, M.S., Lecturers in Orthopaedics
for patting their profound knowledge and practical
experience at my disposal. Their constructive criticism
and valuable options have helped to shape the study to
its present form.

words, but I shall be failing in my duty by not mentioning them. My numerous colleagues - both departmental and otherwise, especially - Dr. A.K. Hai, Dr. S.K. Gupta, Dr. S. S. Dixit, Dr. Ravi Agarwal, Dr. K.K. Pandey, Dr. S. K. Pathak, Dr. Gopal Gupta, Dr. Sunil Gupta and Dr. Vijay Bhandari - all helped me at various stages of this work. Their help is unaccountable.

I am very thankful to ward-boy Satya Prakash who worked day and night for the success of this work.

I am expressing my thanks to Mr. Phool Chandra Sachan who has typed the thesis nicely.

I dedicate this work to the love, understanding and patience of my parents and other family members which has sustained me throughout.

Dateds

Nivajdain.
(Niraj Jain)

CONTENTS

		Page No
1.	INTRODUCTION	1 6
2.	REVIEW OF LITERATURE	7 - 35
3.	MATERIAL AND METHODS	36 - 45
4.	OBSERVATIONS	46 - 61
5.	DISCUSSION	62 - 80
6.	CONCLUSION	81 - 82
7.	BIBLICGRAPHY	83 - 94
	APPENDIX	198
	SUMMARY	(Attached in separate

cover)

INTRODUCTION

Infections pose a great challenge to a clinician from the time immemorial and since the same moment clinicians have been trying their best to wipe out this problem.

thought that the infections could be controlled and infact it had almost done. But there still remains a type of infection against which we are still fighting and several experimental studies have been done in the current century expecially to conquer the battle. That problem is the infection of bones and this challenge further hightens, if the infection is a chronic one, i.e. chronic osteomyelitis.

every year for the treatment of chronic osteomyelitis, but a number of patients return periodically with flairups and recurrent discharging sinuses. One of the difficulties in the treatment of chronic esteomyelitis by chemotherapy is the relative isolation of the infected tissue.

chronic osteomyelitis is more a disease of chronic ischaemia than one of chronic sepsis. Involved areas of bone demonstrate organisms trapped in cancellous spaces by fibrin clot. Pertiens of cortex and marrow may show sequestration. The fibrin clot not

only protects the host from the organisms, but along with the avascularity buffers the organisms from the body's defence mechanisms.

sir Alexander Fleming observed that locally applied chemotherapeutic agents are unable to sterlise a chronically infected wound.

hasmatogenous esteomyelitis present a problem of altered vascular permeability and increased medullary pressure. The problem of ischemia is not encountered until medullary pressure is sufficient to obstruct nutrient artery flow. This apparently requires 3 to 5 days. When treatment is started after the 3rd day, the chance that one is dealing with an early chronic esteomyelitis is increased greatly.

The disease only does not cause agony and discomfort to the patients but also poses as a social problem being a great drain on the human resources and deteriorating the social status by inhibiting the serning capability of already poor class of persons who are main target of the disease.

The problem is further aggravated by paychosocial trauma to the patient due to functional disability caused by parsistant foul smalling discharge from sinuses, constant suffering and chronic bed confinement. The problem is still made worse by the

common occurrence of compound fractures which are most common cause of chronic osteomyelitis.

primary local affection characterised by avascularity because of trauma itself. As this type of esteemyelitis represents both as ischaemic and infective problem since very begining there are fair chances for its conversion into the chronic osteomyelitis. However, provided proper surgical debridement is done, in most of these cases osteomyelitis occurs primarily because of the content—nation of the wound accompanied by less of skin and adjacent soft tissue.

The rapid modernization and industrialisation and road side accidents are the major catalysts behind this scene.

osteomyelitis is full of disappointments. Complete resolution occurs in many other types of infection with the use of antibiotics, but not always with chronic osteomyelitis. This failure must be due to the inherent nature of the bone lesion. Two processes are present: destruction and repair. Destructive processes are seen in the form of cavities containing granulation tissue, pus, organisms, sequestra and fibrous tissue. The reparative process is seen in the formation of new bone, often sclerotic and poorly vascularised. The infective

erganisms are wide spread in this and in the adjacent cavities and sequestra, and these are relatively isolated from the blood circulation and parenterally administered antibioties.

Satisfying experience for the surgeon who understands the basic pathologic process. The primary treatment of any infection is appropriate antibiotics in adequate dosege, coupled with efforts to improve the body's defence mechanisms. Treatment failures are usually due to the inability to deliver adequate doses of antibiotics at site of infection. Dead bone acts as a reservoir for bacterial growth and the dense scar tissue which strounds the infection serve as an impermeable compartment to circulating antibiotics.

Even after removal of dead tissue and sequestra, excision of sinuses and saucerination the dead space left behind is filled up by haematoma which acts as a petential site for the infection. If haematoma and dead space are controlled by open packing, there is epportunity for cross infection.

Elimination of dead space and prevention of cross infection have been essential features of established methods of treatment.

The general principles of previous methods of treatment can be summarised as follows:

- 1. Mobilization of host mechanisms, including rest.
- 2. Radical debridement of involved tissue.
- 3 Elimination of haematoma (dead space).
- 4. Prevention of cress contamination and recontemination.
- The use of antibiotics to decrease mortality and morbidity and to aid local host mechanics.

Continuous irrigation-suction technique incorporates all principles of treatment :

- " With the patient at bed rest, heat defence are mobilized.
- Radical debridement eliminates involved tissue.
- Dead space hasmatoma is controlled both physically by the irrigating solution and by certain possible physiologic mechanisms.
- Primary closure of skin and soft tissues prevents cross contamination and re-infection.
- The presence of antibiotics in the irrigating solution aids local host mechanisms.

This process was first introduced by Duman and Carrel (1917). This method of treatment was first employed in the united states in its current term by Mc Elvenny. Its use in Germany since the mid 1950's has been reported by Willenegger and Roth.

It is the treatment of choice in all cases of established chronic osteomyelitis and all cases of acute hematogenous osteomyelitis warranting medullary decompression.

That is why we performed the present study to evaluate the effect of continuous irrigation-suction technique in the treatment of chronic osteomyelitis by using antibiotic solution.

REVIEW OF LITERATURE

Osteomyelitis is inflammation of all components of bone i.e. inflammation of cortex, meduliary canal and presence of Periostitis.

chronic non-hematogenous osteomyelitis defined as a bone infection that persists for more than six weeks after the initiating episode, with persistently exposed bone, continued drainage, and positive culture of material taken from the bone (May Jr. et al, 1982).

chronic osteomyelitis can be defined as "One or more foci in bone that contain pus, infected granulation tissue, sequestra, a draining sinus, and resistant callulitis" (Andrew J Weiland, MD, J. Russell Moore MD et al. 1984).

The inflammatory foci are surrounded by sclerotic bone with poor blood supply and are covered by a thick relatively avascular periosteum, scarred muscle and subcutaneous tissue. Antibiotics reach such tissue mainly by diffusion. Sensitive organisms may survive and become active again after the therapy is discontinued. Secondary infection with organisms that are more resistant to antibiotics than the primary infecting agent is not uncommon (Fitzgerald Jr., 1983). This changing pathogenesis of chronic esteomyelitis may account for the therapeutic failures that are often seen despite the extensive array

of antibiotics presently available (Fitsgerald Jr., 1979 and O'Riordan, Colm et al, 1972).

The voluminous literature on the subject clearly indicates that none of the methods available for its treatment boday is entirely satisfactory. To begin with the ancient methods of putting seeds in the wound to enhance pus formation, upto present day trials of magnetism and iontophoresis have not shown satisfactory results and the disease is as dreaded as before.

INCIDENCE OF OSTEOMIZLITIS

West et al (1970) said that femoral and tibial osteomyelitis represent 77% of all osteomyelitis seen by the orthopaedic surgeon.

haematogenous osteomyelitis for the year 1951 through 1961 accounted for 44% of esteomyelitis of femer and tibia. Acute haematogenous osteomyelitis is uncommon in adults. However, acute esteomyelitis in drug addicts, especially with involvement of spine, has been reported (Wiesseman et al, 1973).

Occurrence of osteomyelitis with non union fractures in 40% cases described by Kelly. Osteomyelitis is most common in the long bones, particularly lower extremities (Waldvogel et al. 1970 and Edeiken et al. 1973). In neonates, multiple sites of infection are relatively common. Vertibral involvement is next common site in adult (Delinks et al. 1975 end Wald et al. 1980).

The spine is a common site for infection in drug addicts (Kido et al, 1973 and Salahuddin et al, 1973) and in patients with septicemia, including bacterial endocarditis (Mund et al, 1980).

An increased incidence of esteomyelitis occurs in systemic disease, including diabetes (Friedman and Rakow, 1971) and Sickele cell disease (Reynold, 1977).

patients with osteoporosis have an increased incidence of mandible and maxillary esteomyelitis (Gwinn and Free et al, 1972). Patients with insensitivity to pain like diabetes, leprosy also have an increased incidence of osteomyelitis.

PATHOMYSIOLOGY

et al, 1976 and Wald et al, 1980) have divided osteomyelitis into three types that are based upon the variations
in the metaphyseal blood supply in different ages. In
the neonate and the adult, vascular communications are
present between the epiphysis and metaphysis. In childhood (1 to 16 years), the epiphyseal plate separates the
metaphyseal and epiphyseal vascular supply and acts as a
barrier to the spread of infection. The periesteum is
loose in the newborn infant and becomes more tightly
applied to the cortex with increasing age.

In meanates, osteomyelitis frequently spreads to the edjacent joint and produces potentially permanent growth aberration. Extensive periostitis is frequently

present. In childhood, the epiphyseal plate acts as a barrier to infection and joint involvement is less common. Periesteal new bone formation may be extensive. In adults (After closure of epiphysis), joint involvement may occur, but periesteal new bone formation is considerably less prominent.

The medullary canal of the metaphysis is the most common site for hematogenous esteemyelitis. The pus formed within the metaphysis may obstruct the vascular channels and extend into the Maversian systems and Vokmann canals. The avascular infected fragments of dead bone are called "Sequestra" (Waldwood et al, 1970). When the infection extends through the periosteum, exuberant periosteal reaction with new bone formation (involucrum) results. Although hematogenous osteomyelitis is most often seen in childhood, there has been a definite shift towards later caset (Vasly and Waldvogel, 1980). When the metaphysis is within the joint capsule, septic arthritis frequently accompanies osteomyelitis. This is particularly common in the hip or shoulder. Waldvogel found that the incidence of hematogenous osteomyelitis is decreasing and has accounted for only 19% of patients with osteomyelitis seen in the last 10 years. Osteomyelitis may also occur secondary to treuma er inoculation (i.e. drug addicts) or by direct extension from adjacent soft tissue infection. In these patients, infection may occur in any part of any bone.

Osteomyelitis may occur secondary to spread from adjacent soft tissue infection. This is particularly

common in patients with neurologic abnormalities and paraplegia. Osteomyelitis can follow penetrating trauma or open fractures.

Privial trauma has been reported upto 30% of patients with esteemyelitis (Aegarter and Kirkpatrik, 1975 and Waldwogel et al. 1970).

Osteomyelitis can also occur in Pintrak (Stuart A green and Magaret J. Repley, 1984).

Osteomyelitis is also seen after certain surgical intervention like osteomyelitis of Pubis after radical hystrectomy operation (Udo B. Hoyme et al, 1984). Kejlewski et al (1983) reported 5 cases of multifocal chronic osteomyelitis of unknown etiology. In all cases blopsy confirmed the diagnosis but no pathogen was found.

MICROBIOLOGY OF OSTEOMYELITIS

Hematogenous osteomyelitis is usually caused by gram positive organisms, most common being Staphylococcus aureus (Aegerter et al, 1975, Waldvogel and Vasey, 1980).

Kelly (1977) reported that predominant organism cultured was Staphylococcus aureus in about 40% patients of chronic osteomyelitis.

Group B Streptococcus haemolyticus is a common cause of neonatal osteomyelitis. Gram negative organisms, pseudomonas aeruginosa and Klebsiella pneumoniae, have been reported in intravenous drug abusers (Kido et al. 1973).

Gram negative organisms can also be seen in hespital

acquired infection (Grieco, 1972). Other common gram negative organisms include Proteus mirabilis, Enterobacter Sp. and Escherichia colii Kelly, 1977).

The remaining category includes anaerobic organisms such as Corynebacterium diphtheriae or Propionibacterium acnes (Morrey et al. 1977).

albicans can be cultured from drug addicts, patients with diabetes mellitus (Hirschmann and verett, 1976), immuno-suppressed patients on long term antibiotic therapy and premature infants (Svirsky-Fein et al, 1980). There is an increased incidence of salmonella osteomyelitis in patients with Sickle cell disease. Disseminated fungal infactions including coccidioidomycosis (Dalinka, Dennemberg et al, 1971), blastomycosis (Riegler et al, 1974), and actinomycosis may have osseus involvement in addition to pulmonary and cutaneous manifestations.

DIAGNOSIS OF CHRONIC OSTEOMYELITIS

May et al. (1982) described various clinical features of chronic osteomyelitis secondary to trauma or surgery as "a bone infection that persists for more than six weeks after initiating episode, with persistently exposed bone, continued drainage and positive culture of material taken from the bone".

Andrew J. Weiland et al (1984) described clinical features of chronic osteomyelitis in definition as "one or

more foci in bone that contain pus infected granulation tissue, sequestra, a draining sinus, and resistant cellulities. Pus culture and sensitivity is very helpful for the treatment of chronic osteomyelitis in selecting the effective antibiotic.

Erythrocytic sedimentation rate (ESR) which is raised in esteosyelitis is also helpful to assess the course of the disease.

Radiography is very helpful for the diagnosis of chronic osteomyelitis. Capitanio and Kirkpatrik(1976) have described that earliest change in the oesteomyelitis is deep soft tissue swelling which occur within 3 days of the onset of infection. It is characterised by the displacement of the fat lines that are normally parellel to the bony surface. 3-10 days following infection, there is muscular oedema and obliteration of the normally seen thin planes of fat. At the end of this time period superficial subcutaneous cedema is also identified. The earliest osseous manifestations reflect hyperemia and trabecular destruction. Periostitis may be seen as early as 3 or as late as 6 weeks following the onset of infection (Edeiken and Hodes, 1973). Periosteal new bone formation can be lamellated or solid. Solid periosteel new bone fermation is typical of benign lesions (Edeiken and Hodes, 1973). Small periosteal "Dumps' may be seen with granulematous inflammatory disease (Edeiken and Hodes, 1973).

The infection enters the chronic phase as the periosteal new bone proliferates to become the involverum

which eventually completely surrounds the shaft. Heles or skip areas within the involucrum called "Cloaces". Abscess may appear as soft tissue masses obliterating the intervening fat, which differentiate them from tumours, which usually displace the fat planes (Murry and Jacobsen, 1977). Sequestra represent area of dead bone isolated from the blood supply. They can be very small or as large as the entire diaphysis (Murry and Jacobsen, 1977). Sequestra and involucra usually take at least three weeks to appear.

Bone scanning has had a tremendous impact on the radiologic evaluation of osteomyelitis. Positive scans may be seen as early as 24 hours following the onset of infection. The bone scan is uniformly abnormal in chronic osteomyelitis but scan remain abnormal until bone healing and remodelling is complete (Howie et al. 1983). In this situation scanning with Ga⁶⁷ or labelled leucocytes may be helpful in separating such reparative changes from continuing infection (Woolfenden et al. 1980).

wajih Al-Sheikh et al (1985) used In 11, Ge 67 and Tc 99 metadiphosphonate bone scintigraphy for the diagnosis of subscute and chronic bone infections. Scintigraphy was done 24 hours after reinjection of autologous leucocytes labelled with 300-500 u Ci of 111 In omine. Then patients were studied 3 hours after intravenous injection. 20 mCi of Tc 99 MDP and 24-72 hours of Intravenous injection of 5 mCi of 67 Ga - citrate.

to vescular compromise by pus. The cold area often indicates concomitent infection (Berkowitz and Wenzel, 1980). Bene scanning is less accurate in meanatal esteomyelitis. This may be related to the unique anatomy and physiology of the metaphysis in that age group (Ash and Gilday, 1980).

Ai-Scheich et al (1985) described following diagnostic procedure for an orthopaedic patient who is suspected of having chronic bone infection -

Orthopsedic patient with clinical suspician of infection. Radiography Findings diagnostic Negative or Questionable of infection Aspiration and / or treatment for chromic bone infection Bone scintigraphy Positive Negative scintigraphy No infection Incongruent or more Negative or congruent intense lesion as or less intense lesion as compared to Te 99 MDP compared to Te High probability for Low probability for infection, Aspiration infection, loosening and/or treatment for of Devices or other infection. cause of symptoms

Computerised grial tomography (CAT) may be of value in patients with scute esteomyelitis. It can show the extent of medullary involvement even when the plain rountgenograms are normal (Kuhn and Berger, 1979). Soft tissue involvement is well seen on computed tomography(CT).

Vivian W. Wingh et al (1985) studied the CT changes in the 25 patients with acute exacerbation of chronic esteomyelitis. Scan were obtained at 5-10 mm. Sections isolated bony fragments within crotical defects or in adjacent medullary cavity were considered highly suspicious for sequestra.

OSTE OMYELITIS MANAGEMENT

various methods with heavy doses of antibiotics are used every year for the treatment of chronic osteomyslitis, but a number of patients return periodically with
flare-ups and recurrent discharging sinuses, one of the
difficulties in the conventional treatment of chronic
esteomyelitis i.e. sequestrectomy and saucerisation with
chemptherapy is the relative isolation of remaining
injected tissue. Because bacteria may be wide spread in
the Haversian canal or within the medullary cavity.

surgical therapy has decreased the mortality but not the prevalence of osteomyalitis (Waldvogel et al, 1970).

Various methods have been attempted for eradication of the disease. Some workers have carried out this by local application of topical agents in the term of use of het oil (Pere), see sponges (Long, 1927), cambolic acid and

Bichloride of mercury, vaseline gauge, Maggots (Baar, 1931).

Saline (Brockway, 1932) and bacterio-phages (Albee, 1933).

Local chemical sterilization do as much injury to local

host defence as to the bacteria and are ultimately

advantageous to the later (Bryson and Mandell, 1964).

Orr (1927) was the first surgeon who logically applied the principle of rest and immobilisation of the effected part in plaster of paris and augmented it with surgical debridement i.e. sequestrectomy, sewerisation and open packing of resultant cavity. Wound is allowed to heal by secondary intention.

osteomyelitis for many years but this treatment had two major disadvantages - first the healing usually takes a longer time, second the healing takes place by granulation tissue which is later covered by thin epithelium and this epithelium is easily damaged.

ANTIBACTERIAL AGENTS IN THE MANAGEMENT OF OSTEOMYELITIS

Treatment of bone infections with antibiotics is one of the major advancement in the present century. Before the discovery of antibiotics, one third of the patients used to die of acute hemategenous estecmyelitis, and a high proportion of patients with chronic esteemyelitis came to amputation.

Kelly (1977) described a regimen of four weeks of parenteral antibiotic therapy. He claimed the success rate of 90% for chronic osteomyelitis in the categories of

and osteomyelitis in the/mited fractures. But he himself said that it is difficult to prove that this success was attributable to a standardized regimen of antibecterial treatment. Major problem with the antibiotic therapy in the treatment of chronic osteomyelitis is the poor penetration of antibacterial agents in relatively avascular tissue and necrosed tissue or pus.

Though researches claim that Fucidin (Rowling, 1970) and Lincomycin (Magen, 1978) may permeate in the tissue with relatively poor vescular supply and even in the pus. However none of the drug is going to completely cure the condition. Therefore entibiotics are having only supportive role and in no way can replace adequate surgical treatment

SURGICAL MANAGEMENT OF CHRONIC OSTEOMYELITIS

In surgical management of chronic osteomyelitis certain objectives are evident.

First is removal of any dead bone. Sequestra that if remained in a cavity will keep osteomyelitis smoulding for years. Often as a consequence of removal of dead devitalised and infected bone, a sizeable cavity or "dead space" will exist.

second objective is consideration of mathod to obliterate any dead space. A method of obliterating dead space can be a very major aspect of the surgical considerations.

Third objective is to obtain soft tissue coverage of exposed bone.

Fundamentally, the surgical procedure is designed to obtain closure and obliteration of dead space when finely divided bone grafts of cancellous bone are employed to fill a saucerised cavity. The technique is termed as papineau-procedure (Papineau, 1973).

Bone Grafts

12

Lieutenant Coloney Marvin P. Knight et al (1945) described three stage method for the obliteration of bone cavities following traumatic osteomyelitis. The technique was employed extensively in World War II over army personnel injured (Rhinelander, 1975). Out of 23 cases they reported 21 cases have entirely healed wound.

Stage - I: Sequestrectomy is performed and cavity is prepared. Then the wound is kept open with vaseline gauze, and the affected extremity is immobilized.

Stage - II : Five to seven days later split thickness skin grafts applied over the granulation tissue present over the surface.

Stage - III : Obliteration of bone cavity with autogenous bone chips.

Principal factors in the success of this stage are :

 Planning of suitable full thickness covering for the defect.

- Complete excision of the split thickness covering from the depth of the saucerised area.
- Procurement of a sufficient amount of bone chips to fill the cavity.
- Covering of the chip filled cavity with skin and subcutameous tissue in such a manner that tension is avoided.

This method appears promising for the most difficult osteomyelitis of the tibia and femour, the infections involving the ends of a long bone.

Disadvantage of this method is that it requires multiple surgical interventions.

Autoconous Bone Grafts

Ser.

Coleman et al (1946) advocated a single stage procedure of saucerisation, bone grafting and closure with reported success of 92% of 52 cases.

Prigge (1946) reported success in 16 out of 20 cases. He employed grafting with cancellous iliac bone.

Dickel et al (1953) with 36 cases, Hamlette (1954) with 101 cases, Poppirow (1960) with 60 cases, D. Oliviera (1971) with study over series of 120 cases used the autogenous cancellous bone grafts in the treatment of chronic osteomyelitis.

Papineau (1973) and Burri (1975) have reported favourable results using apparently paradoxical technique of grafting free autologous grafts of cancellous bone.

Homografus and Fibrin Adhesion

W. Lack, P. Bosch, and H. Arbes (1987) published their study of 29 bone defects caused by chronic osteomyelitis have been treated by radical excision of mecrotic bone followed by packing the cavities with cancellous homografts held in position by a fibrin scalant. At follow up three years after operation all except one had healed. The removal from homografts of the cellular constituents, such as bone marrow and blood and their subsequent sterilisation by irradiation diminished their antigenicity (Burwell, 1964) and the introduction of fibrin adhesion system improves their recorporation and remodelling (B'each et al. 1976 and B'osch, 1981).

CLOSURE OF OSTEOMYELITIC DEFECTS BY MUSCLE AND MUSCULOCUTANEOUS PLAPS

of muscle flap transposition to cover esteemyelitic defects and Ger (1972) popularized the method and extended the indications to provide coverage for a variety of lesions in the log. In recent years, the use of muscle and musculecutaneous flap transposition to provide vascularized soft tissue cover has resulted in successful treatment of patient with defects resulting from trauma and esteemyelitis (Ger, 1970 and Ger, 1977). James and Gruss (1983) used muscle and musculocutaneous flap for the closure of esteemyelitic defects in 7 patients and reported good result.

Advantages of muscle flap transposition include increased vescularity from locally available muscle which obliterates bone cavities and fill soft tissue defects, preventing leaking of emudate, allowing antibiotic to perfuse in infected area, permitting ready up take of split skin graft, and preventing afherence of skin to bone (Oct. 1977).

PREE TISSUE TRANSFER

夢

microsurgery have made it possible to transfer free cutameous, myocutameous, muscle, bone and entercutameous flaps from one part of body to a recipient site that is deficient of such tissue. Blood supply is restored to the free graft at the transfer site by microvascular anestameous of donor and recipient arteries and vains. This method predictably results in healing of esteemyelitis (Cordon and Buncke, 1983; Nathews et al., 1982 and May et al., 1982).

May Mr. and Savage (1963) stated in their report
that estecmyelitis should be treated aggressively by a
reconstructive microsurgeon. In 1982 May et al reported
their results with microvescular transfer of free tissue in
22 patients with 100% success. The series was later
expanded to 30 patients with only one failure.

Mathews et al (1982) described the use of free muscle flaps in 11 patients with chronic esteemyelitis with me requirence of infection.

Gordon and Bunke (1983) described the results in 14 patients with esteomyelitis who had been treated with repeated debridements and free flap coverage. Two patients had recurrence of the infection.

Andrew, J. Weiland et al (1984) described 79% success in the 37 free tissue transfer.

ADJUVANT HYPERBARIC CHYGEN

pavis, J.C. recently reported (in press) the results of a follow up evaluation of 40 petients of chromic ostoomyelitis who were treated with systemic hyperbaric oxygen as an adjuvant to debridement of the infectious focus in bone, as well as systemic administration of antibiotics. He reported that 75% of the original patients remained free of infection.

Morrey et al (1979) used hyperbaric oxygen in the treatment of chronic osteomyelitis. He published the series of 40 patients. Jefferson C davis et al (1986) described the use of adjuvant hyperbaric oxygen in the treatment of chronic osteomyelitis in 38 patients. They reported 34 patients with successful results. Patients were treated by local debridement of the wound, prolonged parenteral administration of antibiotics and once a day treatment with hyperbaric exygen, for on an average 48 days.

SILVER IONFOPHORESIS

Use of electrically generated silver ions in treating chronic osteomyelitis was described by Becker

et al (1978), and Webster (1981). It has got both bactericidal as well as osteogenic properties.

TOPICAL ANTIBIOTIC - DETERGENT TECHNIQUE

this method in cases of chronic esteemyelitis. They used Penicillin in sodium chloride solution along with wetting agent 'Tergitol'. He emphasized the non-eparative treatment of wound, while Mitra used this technique after wound debridement. The one better aspect of this method is that it is simple and so can be widely used in these areas who are underdeveloped and without facilities of extensive surgery.

CONTINUOUS IRRIGATION AND SUCTION TECHNIQUE

publica.

The treatment of chronic esteemyelitis is very cumbersome, time consuming and unsatisfactory till date which signifies from the fact that several attempts to cure it, have been made starting from boiling water to various reconstructive operative procedures of today, but none is fool proof.

Now the question arises that why the chromic osteomyelitis is notorious one. First and the most important the antibiotics could not reach at the site of infection. Thereby persistence of bacteria in that part of bone which is rendered avascular by them. Second, antibiotics could not transgress the area occupied by necrosed tissue and pust third there occurs the thrombosis of subperiosteal vessels

and main nutrient artery thus blocking the way of antibiotics to infected part of bone.

Even after removal of dead tissue and sequestra excision of sinuses and sequestration, the dead space left behind is filled up by the heematoma providing a potential site for the infection.

in the cavity thus formed, the infection occurring subsequently will not take place and this brought the idea of the technique of continuous irrigation and suction. Doing irrigation by antibiotic solution the additional advantage would be the elimination of bacteria, residing in Hayersian system and medullary cenal.

This process was first practised by Duman and Carrel (1917).

annulse" were made for the osteomyelitis service of the Massachusells General Hospital. These were stainght glass canulae, each with a flange at one end to prevent it from slipping out of the wound. After incision and drainage of an osteomyelitic area two of these cannulae inserted and at regular intervals irrigation fluid was allowed to flow through the wound. Glass was not ideal because of possibility of breakage, so in 1938; the vitallium cannulae were made. Smith Petersen (1945) used such cannulae for drainage and irrigation in the treatment of septic wounds by local chemotherapy with primary closure. They used Dakin's solution, silver - pectinate and Penicillin as local chemotherapeutic agents in irrigation fluids. Duration

of irrigation was 2 to 4 weeks. When irrigation was complete, cannular removed under pentothal anaesthesia and soft tissue defects were closed. They also used systemic chemotherapeutic agents along the local chemotherapeutic agents.

In 1942 Brantigan and Owen reported the topical use of sodium tetradecyl sulphate in the treatment of acute pyogenic empyema of pleural cavity. This acts as detergent or "welling agent".

topical use (in 3 cases of chronic esteomyelitis) of penicillin in a solution of isotomic sodium chloride with 0.1% of the detergent sodium tetradecyl sulphate.

results following the topical use of a penicillin detergent solution in treating 2 was veterans who had been discharged from millitary service with intractable osteomyelitis.

In this same report they referred to their success in treating a series of 37 civilian patients who were suffering from chronic esteomyelitis.

During the decade following his first successful use of a detergent antibiotic solution, Grace and his co-workers were able to effect a cure of many cases of infection in vareity of bacterial organisms including the tubercle bacillus. Refinements in the technique included the closure of the wound before beginning instillation of the detergent antibiotic solution. Antibiotics which were soluble, in the solution and also previously determined

114

of Calcutta, India worked with Grace in Brooklyn, and in 1957 they reported compete arrest of 64.21% of 95 cases of chronic pyogenic esteemyelitis treated in Calcutta and 69% of the 45 similar cases treated at the Grace Clinic, Brooklyn, New York. In these cases the actual surgery performed consisted only of removal of sequestra and other grossly devitalized tissue and opening the medullary canal sufficiently to assure free entry of the detargent antibiotic solution. They used "Aerosel Wash" solution as detargent.

Edward L. Compere (1962) used above mentioned technique in the treatment of osteomyelitis and infected wounds by closed irrigation with a detergent antibiotic solution. He used 5 cc of the detergent antibiotic mixture. Instilled through the T-tube 3 times each day for 5 to 10 days. He reported that results are very gratifying which have been echieved in a small series.

containing antibiotics was first used for patients on the orthopsedic service of Chicago Wesley Mamorial Mespital Goldman, Johnson and Grossberg (1960): resident surgeons in orthopsedic surgery described the technique and reported five cases of osteomyelitis which were cured by continuous or intermittent closed irrigation. The time elapsed between beginning circulation and healing of the wounds was from 1 to 8 weeks.

Mc Elvenny (1961) developed the detail technique described by Gedman, Johnson and Grossberg. He reported 12 cases treated by this method. How ever its use in Germany since mid 1950's has been reported by Willenegger and Roth.

Dombrowski and Dunn (1965) reported the comparative study of the short term results of closed wound irrigation and suction technique with those obtained by the open packing method in another series of cases. They carried out 29 closed irrigation procedure of major bones in 22 patients. Results of these cases were compared with 16 open packing procedures. They claim that failure rate with open packing (43%) were higher than with closed irrigation (23%). More over they also mentioned that closed irrigation suction incorporated all previous principles of treatments ; with the patient at bed rest, host defenses are mobilized. Radical debridement eliminates involved tissue. Dead space hematoma is controlled both physically by irrigating solution and by certain possible physiologic mechanisms. Primary clesure of skin and soft tissues prevents cross contamination and recontamination. The presence of antibiotics in the irrigating solution aids local host mechanisms.

Dombrowski and Dunn (1965) postulated the following explanation of why prolonged irrigation may seem to substitute atleast in part even for inadequate debridement. There is an osmotic gradient of 15 m of Hg. between irrigation solution and interestitial fluid.

This includes a flow of fluid from tissue to wound cavity.

Second factor which helps the flow of fluid is temperature because temperature of tissue is about 37°C and that of irrigation fluid is about 25°C. This flow brings fresh lymph and leucocytes into the area which have antibacterial property and help in obliteration of dead space.

method for closed system, irrigation-suction, antibiotic, detergent treatment of deep wounds. Nost of the 24 petients in series had acute infections of the muscule-skeletal system and there were some chronic infections.

many of the time-honoured measures such as sequestrectomy and debridement, but particularly an effective closed irrigation-suction tube system. The use of clerpatic -NCB and a wide selection of antibiotics, whose local effect appears to be increased when they are combined with detergents (Alevaire) (Grace and Bryson, 1945; Mitra and Grade, 1956) directed with considerable precision to the involved parts, makes unnecessary and underireble wide spread removal of bone essential for structural stability. The suggested flow reversal technique to avoid plugging of tubes.

They claim the results as gratifying and severe residual crippling due to tuberculosis of hip joint in children may be prevented by this method.

Polo (1968) employed the irrigation-suction antibiotic detergent method in 8 children with tuberculosis of hip and 4 children with tuberculosis of knee and claimed favourable results.

in the treatment of acute heamstogenous esteomyelitis with marked success. They claimed that duration of this type of management could be reduced to only a few days

closed instillation-suction for the treatment of chronic esteomyelitis in 12 patients. In their technique, infected bone is first exposed and all necrotic material removed. Three perforated drainage tubes are inserted. Closed continuous-instillation-suction thus established, and had been maintained for upto 6 weeks. They claim results as follows:

- . In one patient method failed technically.
- In 8 cases out of 11 in which the infection was resistant to conventional treatment. There was clinical resolution of the infection with the instillation suction technique.

Lawyer et al (1972) described intermittent closed irrigation technique for treatment of osteomyelitis.

Michelinakis (1972) published the result of continuous-irrigation-suction method used for the treatment of chronic osteomyelitis in 12 patients. The irrigation solution he used, composed of 800 ec normal saline, 200 ml Alevaire and 2 antibiotic according to the culture and sensitivity reports. He used Y-shaped tubes and one of the each Y tubes lies within the medullary cavity and other in the surrounding soft tissue. He described good results in 10 patients and they remained clinically symptom-free and there was no radiological evidence of infection for 2 to 4 years.

He also mentioned that all the patients had previously failed to respond to conservative treatment with antibiotics, and immobilization etc. and 8 of them had from 1 to 7 operations for the infection prior to irrigation (i.e. sequestrectomy, excavation of abscess etc).

Clowson, Davis and Hansen (1973) published their study based on treatment of chronic osteomyelitis with emphasis on closed suction-irrigation technique. Characteristic features of their work were :

- i. They used different methods for the treatment of chronic esteemyelitis in 176 patients and results of treatment with suction-irrigation technique compared with patients treated by other types of surgery.

 The results had been considered by the 3 parameters namely drainage, pain and toxicity.
 - They mentioned the following important technical considerations in the use of suction-irrigation -

- a. They placed tubes parallel, side by side to each other and tied both tubes to each other.
- b. They recommended the irrigation flow rate of 1 to 2 litres per hour for the first 2 days.
- c. To avoid frequent change of bottles they used renal dialysis unit.
- d. They performed irrigation in 5 to 40 days.
- e. They also gave emphasis over the good nursing and after care.
- f. Initially detergent and antibiotics were used in the irrigation solution but later on they did not used them.
- 3. Results were published as follows :
 - a. Out of the patients treated with the suction irrigation method, 73.5% were asymptomatic at follow up as compared with 45% of those treated by other methods.
 - b. 80% of the patients treated with suctionirrigation were clinically improved by atleast one parameter as compared with 78% improvement in patients treated by other methods.
 - c. 47 patients underwent girdle stone Hip orthoplasties 18 were treated epen and 29 were treated
 by closed suction-irrigation method. Of the 18
 treated without suction irrigation, only 8% were
 asymptomatic at follow up and only 69% were
 clinically improved by at least one parameter.

Of the 29 patients treated with suction irrigation, 74% were asymptomatic at follow up and 97% were improved clinically by atleast one parameter.

Letts and Wong (1975) treated 18 cases of acute ostsomyelitis with closed tube irrigation and their results were compared with a similar group in which the treatment was the same except for the omission of closed tube irrigation (Incision and drainage etc.).

They mentioned that technical complications were developed in 13 of the 18 patients, in whom cleans tube irrigation was used. Post operative hospitalization was almost double in case of closed tube irrigation than that of treated with other method.

They concluded that the end result of treatment of acute osteomyelitis with closed tube irrigation is little different from that of simple incision and drainage and antibiotics. And suction irrigation in the treatment of acute osteomyelitis in children does not appear to be essential for successful management and introduced added risks of superinfection and technical complications.

Compare, Metzgeg and Mitra described treatment of pyogenic bone and joint infections by closed irrigation (circulation) with nontoxic detergent and one or more antibiotics.

Rao and Sahu (1973) compared the results of 18 patients of chronic osteomyelitis treated with suction methods like : sequestrectomy, saucerization or both. The irrigation fluid was normal saline with the appropriate antibietic. Of the patients treated with suction—irrigation technique 77.7% were asymptomatic at follow up as compared with 43.3% of those treated by other methods. Poer results from the use of closed suction irrigation is usually from one or more causes as stated by Clowson et al (1965—1973). Inability to remove all the dead bone and surrounding scar tissue, failure to understand the principles of suction—irrigation system and technical failures of surgical and nursing team.

treatment of chronic esteomyelitis with continuous irrigation suction technique in 30 cases. In this series majority of cases had involvement of long bones. They used irrigation fluid normal saline and one or two antibiotics according to culture and sensitivity reports. They used 1000-1500 ml of irrigation fluid in 24 hours duration of irrigation ranged from 4 to 14 days.

The most common causative organism was staphylococcus aureus. It may present alone or in combination with other organisms in 75% of the patients with positive culture.

Similarly, Staphylococcus as a major cause of osteomyelitis had been reported by other weskers (Mitra

and Grace, 1956; Mercer, 1959; Michelinaki, 1972; and Rao and Sahu, 1978). Bajaj et al evaluated the results as "excellent" when there was healing by primary intention, no sinus or residual infection, no pain or local tenderness "good" when there was delayed wound healing or occasional pain and tenderness but no residual infection or sinus; and "Poor" when there was parsistent pain or discharge.

The results of their series were excellent in seven, good in 15(together 73.3%) and poor in 8(26.7%).

irrigation technique as : almost complete prediction that
the operative wound will heal by primary intention—
infrequency of dressing, reduction in the time for
complete healing, minimum loss of serum, no soiling of
bed linen, comfort and lack of systemic reaction,
reduction in cross infection, hospitalization and post
operative isolation. Metalic intramedullary fixation
may be left in place or inserted at the time of operation.

MARRIAL AND METHODS

10

In the present study, a concerted effort has been made to evaluate the effect of continuous irrigation and suction technique by using antibiotic solution in the treatment of chronic osteomyelitis.

clitis of different bones coming from peripheral areas of Jhansi and other adjoining districts were included in this study. These were admitted in the department of Orthopsedics, Meharani Lammi Bai Medical College, Hospital, Jhansi (U.P.) where study was carried out.

CRITERIA FOR THE SELECTION OF CASES

· Jane

- Patients having pus discharging sinus with radiographic findings suggestive of chronic osteomyelitis.
- "Presh cases as well as cases of chronic esteomyelitis treated previously by other methods (whether conservative and / or operative like incision and drainage sequestrectomy, saucerisation etc) have been included in present study.
- Patients of chromic osteomyelitis without sequestrum.
- Patients of chronic osteomyelitis where saucerisation is not possible due to lack of strength of the bone.

Patients not turning up after operation or having fellow up period of less than 2 months were not included in this study.

PRE-OPERATIVE EVALUATION AND PREPARATION

All the cases were thoroughly interrogated for detailed history, examined clinically and properly investigated before any surgical intervention. Probing of the sinus remained an essential step in the clinical examination.

Various investigations done were as follows :

- 1. Blood routine : HbK, TLC, DLC, ESR,
- 2. Urine routine : Albumin, Sugar, Microscopic examination.
- 3. Blood Sugar : Pasting and postprandial or random.
- 4. Blood urea.
- 5. Pus culture and sensitivity.
- Antero-paterior and lateral skiegrams of affected part.

Pre-operative tetanus toomid and entitetanus serum were given in all cases. Systemic antibiotics were given pre-operatively according to the culture and sensitivity report and if pus culture and sensitivity could not be done then crystalline peniceilline and Gentamicin were given.

In general debilitated patients their general condition was improved by oral or parentral iron therapy or by blood administration along with protein rich diet and vitamin preparations. For the selected cases blood was also arranged for per-sperative

transfusion. Only then operations were carried out when patient become fit for anaesthesia.

Proper pre-operative preparation of the part was started well in advance. Twenty four hours before the operation methyline blue dye was injected in sinus, to detect the exact sinus tract per-operatively in selected cases.

studied along with clinical exemination to assess which procedure is to be carried out viz. dril holes, sauserisation, sequestrectomy, syringectomy etc. along with irrigation and suction.

OPERATIVE TECHNIQUES

Under appropriate anaesthesia (either general or regional) operation was performed as single stage procedure under strict aseptic conditions.

- ... Part was first painted and drapped.
- With the help of a probe direction and depth of sinus was determined.
- Suitable incision was made considering the condition of skin, presence of scar tissue and site of sinus.
- Diseased bone was reached by appropriate approach and widely exposed.
- Assessment of condition of bone and surrounding soft tissues was made.
- All fibrosed and necrosed soft tissue, sinus, tract, unhealthy granulation tissue and sequestrum present

were removed.

- Saucerization of the cavity if present was done.
- Medullary canal on both sides was opened.
- Per-operative pus specimen was collected and sent for culture and sensitivity.
- The cavity present was curetted.
- Then wound was thoroughly irrigated with copious amount of normal saline to remove all the debris and blood clots.
- In the cases where saucerisation was not possible and sequestrum was not present two holes were made, one at each end of the bone, through separate incisions, by the help of bone awl or bone dril. Through these holes irrigation tubes were placed inside the marrow cavity.
- Irrigation and suction tubes were placed parellel side-by-side and exiting in apposite direction.
- The tubes were brought out obliquely from the wound through as much soft tissue as possible, penetrating the skin with the trochar atleast 1 inch away from the incision line. One point was kept in mind while placing the tube that perforated part of the tube must not be in the muscle plane and extra part of the perforated tube was cut.
- Tubes were fixed externally with the skin tightly so that tubes must be secure to the skin up to desired period.

- ... Wound was closed in layers and properly dressed.
- Post-operative immobilisation was given either in form of plaster of paris slab or in form of skin traction over Thomas' splint according to the bone involved.
- Irrigation tube was connected to the irrigation bottle containing normal saline and one or two antibiotics.
- .. Suction tube was connected to the suction apparatus.
- Irrigation-suction mechanism started and its functioning checked.

Irrigation Fluid

Normal saline was used for the purpose of irrigation.

SELECTION OF ANTIBIOTICS

- ... One or two antibiotics were mixed in the normal saline according to culture and sensitivity report.
- When culture and sensitivity can not be possible or sterile then crystalline penicilline and Gentamicin were used till report obtained.
- when pus found to be resistant to all antibiotic them also erystalline penicalline and Gentamicin were used.
- During pre-operative and post-operative periods systemic antibiotics were also given.
- Dilution of antibiotics in normal saline was :

 Crystalline penicilline : 10 Lec IU/Lit.

 Streptomycin : 2 gm/Lit.

 Gentamicin : 30 mg/540 ml.

Chloramphenical s 1 gm/Lit.

Ampicillin

a 1 cm/Lit.

Caphasoline

1

s 1 gm/Lit.

- The outcoming fluid was cultured regularly and antibiotics were changed according to the sensitivity report.

MAINTENANCE OF CONTINUOUS IRRIGATION-SUCTION TECHNIQUE

- For the first two days 2000-2500 ml of irrigation fluid per day was used. This is because the problem of blockage in suction tube is maximum during first two days and by keeping the irrigation flow fast this problem was solved.
- Problem of blockage in the tube was solved by reversing the flew of tubes.
- . From the third day enward 1000-1500 ml irrigation fluid was used per day.

DURATION OF CONTINUOUS IRRIGATION-SUCTION TECHNIQUE

- Outcoming fluid was cultured at regular intervals and irrigation was continued till two consecutive cultures of out coming fluid were sterile.
- This technique was usually continued for 2-3 weeks.
- First, irrigation tube was taken out and suction was continued till fluid came out and then suction tube was also removed.

IRRIGATION-SUCTION APPARATUS

- For irrigation normal saline bottles containing 540 ml normal saline were used and appropriate amount of antibiotics was added.
- was connected to irrigation tube.
- Irrigation-suction tubes : Gamma irradiated sterilized p.V.C. tubes of F.G.=10 size were used. These tubes were reasonably stiff because soft tube might colapse under negative pressure. These tubes were having multiple side holes at proximal end.
- Suction apparatus Three types of suction
 apparatus were used :
 - 1. Glass bottle with PVC tubings negative pressure was generated in the bottle by the help of suction machine and this bottle was connected to suction tube. Outcoming fluid was collected in this bottle.
 - Plastic bag with PVC connections megative pressure was generated inside the bag by compressing it and this device was attached to suction tube and outcoming fluid was collected in the bag.
 - 3. 3. Electrical slow suction apparatus Suction tube
 was attached to this apparatus. Advantage of
 this apparatus is that suction pressure can be
 regulated according to the requirements.

- Stitches were removed after 12-14 days depending upon the condition of local area.
- Check skiagrams were taken with the tubes in position to assess the success of operation or any left over piece of sequestrum.
- Later on patients were discharged with the immobilisation according to site and extent of involvement
 and with instructions to continue the prescribed
 antibiotics in appropriate doses for further four to
 six weeks.

POLLOW UP

hospital periodically and were assessed clinically for pain, tenderness, discharge, soft tissue healing or any other complications and radiologically to see the condition of bone whether radiologically there is any persisting evidence of infection(e.g. sequestrum formation).

Type of secondary treatment if given in any case of recurrence or having complications was also noted.

Different workers have laid down different criteria for assessment of results of continuous irrigation and suction technique. But none of the previous workers have laid down the clear cut criteria for the assessment of result. E. Michelinakis (1972)

used following criteria for good result. (a) Duration of clinically symptom free periods and (b) no radiological evidence of infection.

Clawsen et al (1973) considered the results by following 3 parameters : Discharge, pain and toxicity.

In this series definite clinical and radiological criteria were used for the assessment of the results in three grades i.e. Excellent, good and poor.

A. Clinical Criteria

- i. Healing by primary or secondary intension.
- 11. Sinus.
- 111. Residual infection.
- iv. Pain.
 - v. Tendermess.
 - Wi. Discharge.

B. Radiological Criteria

- i. Evidence of infection.
- ii. Sequestrum formation.

GRADES

Excellent

When there was healing by primary intention no sinus or residual infection, no pain or local tenderness no radiological evidence of infection.

Good


when there was delayed wound healing or occasional pain and tenderness but no discharge or sinus and no radiological evidence of infection.

Poor

100

when there was persistant pain or discharge or radiological evidence of infection or formation of sequestrum.

All the above information regarding clinical examination, investigations, operative procedures and follow up findings were noted down in a proforma. The sample of which is attached in appendix.




A : N. SALINE + ANTIBIOTIC

B: IRRIGATION-TUBE


C : SUCTION - TUBE

D : SUCTION - APPARATUS.


DIAGRAM SHOWING IRRIGATION - SUCTION SYSTEM.




Ph. No. 1 : Showing sinus an posterior aspect of thigh.




Ph. No. 2 : Thigh with part painted and draped.



Ph. No. 3: Showing sequestrum through the pro-existing window in the femur.




Ph. No. 4 : Sequestrum was being taken out after increasing the window size.




Ph. No. 5 : Saucerized femur along with sequestrum.


عالمات



Ph. No. 6 : Ingoing irrigation tube inside saucerized cavity.




Ph. No. 7 : Outcoming suction tube inside saucerized cavity.



Ph. No. 8 : Showing stitch line and anchored tubes.

Ph. No. 9 : Showing irrigation-suction mechanism in functioning.

禮



Ph. No. 10: Showing all 3 types of suction apparatus along with irrigation and suction tubes (From L4 to Rt App. No. 2,1, 3).

OBSERVATIONS SECUCIONOS SECUCIONAS SECUCIONA

The present study "To evaluate the effect of continuous irrigation-suction technique by using antibiotic solution in the treatment of chronic osteomyelitis" was carried out on 24 patients of proved chronic osteomyelitis of different bones, admitted in orthopaedic ward, of M.L.B. Hedical College, Hospital, Jhansi between July, 1987 and June, 1988. Patients were treated by continuous irrigation and suction technique by using antibiotic selution. Out of 24 patients 1 patient died post-operatively due to flaring up of septicemia and bacterimia.

Out of 24 patients mode of spread haematogenous in 21 cases (87.50%) and in rest 3 (12.50%) nonhaematogenous osteomyelitis was found (Table I).

Table	1	1	Mode	of	spread	02	osteomyelitis.
-------	---	---	------	----	--------	----	----------------

pread	of osteomyelitis	No.of cases	Percentage
laemato	ogenous	21	87.50
"on-ha	ematogenous		
Provide (M)	In simple fracture	2447	12,50
5	In compound fracture	440	460)
	70t8l	24	100.00

The patients of present series were from different age groups ranging from 7 years to 50 years. Mean age of the patients was 19.79 years. 54.17% patients were from

second decade of life followed by 16.67% and 12.50% patients from third and first decade respectively (Table II).

Table II : Incidence of age distribution of the cases.

(Years)	No. of cases	
0 - 10	3	13.50
11 - 20	13	54.17
21 - 30	4	16.67
31 - 40	2	8.33
41 - 50	2	8.33
750	400	ab
Total	24	100.00


In this study, out of 24 cases, 21 (87.50%) cases were male and 3 (12.50%) cases were female.

(Table III, Fig. 1).

Table III : Sex distribution of the eases.

	No. of cases	Percentage
Sex	NO. Of Conce	
The second secon		A. 579 BY CO.
Malo	21	87.50
Mara		12.50
remaio	3	***
T. CLIVENING AND		
	24	100.00
Total		

According to the socio-economic status of patients, 12 (50.0%) cases were poor, 6(25.0%) cases were from lower middle class and rest 6(25.0%) cases were from middle class (Table IV).






FIG. 1 SEX - DISTRIBUTION

Table IV: Incidence of socio-economic status of cases.

Socio-economic Ftatus	No.of Cases	Fergest 836
Lower	12	50.00
Lower middle	6	25.00
Middle	6	25,00
High	49	400-
$r_{0}=c_{0}+c_{1}+c_{2}+c_{3}+c_{4}+c_{5$	as we represent the contribution of the section of the contribution of the contributi	100.00

radius 3 (12.50%) cases each, Ilium was involved in 1 (4.17%) case only (Table V).

Table V : Incidence of bone involvement in cases

Bone involvement	No. of cases	Percentage
Persur	11	45.83
Tibia	6	25.00
Humorus	. 3	12.50
Radius	3	12.50
Ilium	1	4.17
and the second distribution and the contract of the second distribution $70ta ilde{1}$	we can be a considered an analysis of the constant of the con	100.00

Out of 24 cases, bones of lower extremities were involved in 18(75.0%) cases while bones of upper extremities were involved in 6 (25.0%) cases (Table VI).

Table VI : Incidence of limb involvement.

L1mb	No.	of	00000	
U ppet		6		25.00
Lower		18		75.00
non-turnament sectorement respectively.		24		100.00

Duration of illness in cases of present study
was quite variable ranging from 1/2 months to 8 years. The
average duration of involvement was 17.88 months. Nine
cases (37.50%) were having duration of involvement from
0-6 months. Five cases (20.83%) each were having duration
of involvement from 7-12 months and 13-18 months. One case
(4.17%) was having duration of illness from 19-24 months
while 4 (16.67%) cases were having illness of more than
24 months duration (Table VII).

Table VII: Incidence of duration of illness.

wration (months)	No. of cases	Percentage
	9	37.50
7 - 12	5	20.83
13 - 18	5	20.83
19 - 24	1	4.17
7 24	4	16.67
Total	centres contacumenta con estra concessor in consensor acres con estra transition con consensor con consensor e conte com 2.4	100.00

Involvements of joints was detected by seeing movement arround neighbouring joints when movements found to be restricted then a joint is said to be involved. Joint were involved in 11 (45.83%) patients while rest 13(54.17%)

cases were not having joint involvement (Table VIXI).

Table VIII: Incidence of joint involvement.

Joint involvement	No. c	£ 0080	Porcentage
Present	1	.1	45.83
Absent	3	.3	54.17
To ϵa	1	34	100.00

In present study 23 (95.83%) cases were having one or more discharging sinus while only 1 (4.17%) case was devoid of discharging sinus (Table IX).

Table IX : Incidence of existing sinuses.

An admitted on comment			active over the delication	er-ganglybrennedestandsfreihilderenne	
Discharging	inuses	No.	of	Cases	Percentage
The second districts of the second se				no do	95.83
Present				23	20 4 60 4
1.7 000 0000				-	4.3.7
Absent				1	40 4 3. 1
AND STREET					
And the second s			45(4991749	A 4	100.00
Tota	1			24	
4,000, 100, 100, 100, 100, 100, 100, 100		NEW THE PERSON NAMED IN COLUMN TWO	gratuu kalitähinen 415	to with the control of the control o	

Twelve cases (50%) were previously treated with chemotherapy while 17 (70.83%) cases had one or more incision and drainage procedure and in 5 (20.83%) cases
saucerization and/or sequestrectomy was performed previously (rable X).

Table X : Showing previous treatment.

Type of previous treatment	No. of cases	Percentage
After the second	12	\$0.00
Chemotherapy	2.7	70.83
Incision and drainage Saucerization and/or	5	20.83
sequestrectomy		

In 6(25.0%) cases of present series no surgical procedure was performed before coming to this hospital. In 11 (45.83%) cases single surgical procedure was performed previously while in 4(16.67%) patients 2 surgical procedures had taken place. Three, four and more than four procedures were performed in 1 (4.17%) case each (Table XI).

Table XI: Number of previous surgical procedure.

No. of surgical Procedure	No. of cases	Percentage
0	6	25.00
2	11	45.88
2	4	16.67
3	1	4.17
4	1.	4.17
74	1	4.17
rotal		100.00

following microbiology. Staphylococcus aureus was the commonest organism isolated from 10(41.65%) cases while from 1 (4.17%) case it was isolated along with proteus. Pseudomonas was isolated from 3 (12.50%) cases whereas E.Coli, Klebsiella and Proteus were isolated from 1(4.17%) cases each. E. Coli along with proteus was isolated from one (4.17%). No organism was isolated from 6(25.0%) cases i.e. their cultures were sterile (Table XII).

Table XII: Incidence of various organisms isolated from patients pre-operatively.

organisms	NO.05	Percentage		
staph. aureus	10	41.65		
Staph. aureus + Proteus	1	4.17		
Pseudomonas	3			
Klebsielle	2.	4.17		
E. Coli	3.	4.17		
Protess	1	4.17		
E. Coli + Proteus	1	4.17		
Sterile	6	25.00		
stance and the attack and the control of the contro	24	100.00		

Incidence of super infection was detected by culture of drainage fluid. If organisms cultured were other than the pre-operative organisms then it mean that super infection had taken place. Eight cases (33.33%) in showed super infection while/16(66.67%) cases there was no superinfection.

Pseudomonas was the commonest organism responsible for super infection isolated in 4(16.67%) cases (Preoperative organism was Staph. aureus in 3 cases while in 1 case no organism was isolated).

In 2 (8.33%) cases Protous was the organism

(Pre-operative organism was Staph. aureus in 1 case and

*labsiella in other one), while in 1 (4.17%) case Staph.

aureus was responsible for super infection (No organism was
isolated pre-operatively). In 1 (4.17%) case mixed super

infection had occurred where . Coli along with Proteum was isolated from drainage fluid (Pre-operative organism was Staph. aureus) (Table XIII, Fig. 2).

Table XIII: Incidence of bacterial super-infection in drainage fluid.

Super infecting organism	No.of cases				Percentage					
$Q_{Z}^{*}CCCLA + ODE$	(Marie Control of the	-	NAME OF THE OWN OF THE OWN OF THE OWN OF THE OWN OWN OF THE OWN	general services		4	42	67		
Pagudononas		4				A	13 p	01		
(Pre-Op. Staph. aureus - 3 Sterile - 1)					White a	of Challeton	mine	curron material	- Option (S	
entato diades trancio sociali entato sociale intrino megida algori mondo contro sociale algori titolio entato entato	energia excepto	displa 46	and invest	specials.	- Bouletters	alestors.	Green.			
Proteus		2					8.	, 32		
(Pre-op. Staph. aureus - 1 Klebsiella - 1)										
which wides allow value from these states about more states about states office which which	streets soldier	witers 4	sistile stalle	1900/8	THE STATE OF	equit	ataless .	el selo	in spiller	
Staph. aureus (Pre-op. Sterile - 1)		1					94	,17		
games record through the control and the second states are second states are second states are second states and the second states are second	Jestifia attation	sitesfin i	MANUEL ACCORD	16/10/10	-90000	raphopse	eston	nichese vising	go 446co	
E. Coli + Proteus		1					4.	.17		
(Pre-op. Staph. aureus - 1)	e annie farie	entolis	diller desir	s carrella	s dkilnib	cocide	Spirit	epitino del	(refo (shisy)	
Total incidence of superinfecti	.OD.	8					II.	. 33		
COMPON TOTALD ANGLES SERVING ARROWS A	ale rigorates expensive	4000	assemble streng	is liquidit	p -soluted	wints	ando dis elli	40 CT 10	域 对动	
No super infection		16				ng/squipermi		.67	neppi SVI neppi	
Total		24				1	00	.00	ed and Shift at Mark	

Radiological features present in patients of this study were as follows. Periosteal reaction was present in 23 cases. Sequestrum was seen in 14 cases. Bone destruction was visualized in 21 cases. Sclerosis was evident in 13 cases. Thickening of the bone was noticed in 16 cases while cavity was identified in 17 cases (Table XIV, Fig. 3).




FIG. 2-INCIDENCE OF BACTERIAL - SUPERINFECTION

Table XIV: Showing radiological features.

Padiological features	No. of cases
Periosteal reaction	23
Sequestrum	1.4
Destruction	21
Sclerosis	13
Thickening	16
Cavity	2.7

Various types of operation were performed in different patients along with the use of continuous irrigation and suction technique. Curettage was done in 2 cases, syringectomy was performed in 14 cases and in 2 cases drill holes were made at the ends of bones.

Saucerization was performed in 18 cases while sequestrectomy was done in 13 cases (Table XV).

Table XV: Showing the types of operation done.

Types of procedure	No. qf cases
Curettage	9
Syringectomy	14
Drill hole at the ends of bone	2
Saucerization	1.0
Sequestrectomy	1.3

In the present series healing occurred by primary and secondary intention in 16 (69.56%) and 7(30.44%) cases respectively (Table XVI).




FIG. 3 RADIOLOGICAL - FEATURES.

Table XVI: Showing type of wound healing.

Type of wound healing	No.of cases	Percentage
Primary intention	16	69.56
Secondary intention	7	30.44
Total	23	100,00

Post-operative hospital stay of the patients ranged from 14 to 40 days with the average stay of 22.26 days. Fourteen (60.87%) cases remained in the hospital from 0-3 weeks postoperatively. Seven cases (30.44%) had post-operative hospital stay from 3-4 weeks while 2(8.76%) patients had more than 4 weeks stay (Table XVII).

Table XVII: Duration of post-operative hospital stay.

Dui	cat	ion	(weeks)	No. o	f cases	Percentage
0	100	3			14	60.87
3	490000	4			7	30.43
7	4				2	8.70
cipii galentii de	ndedineda-ipa	To	rtal.	die Addres des reiter der der sonform der mit der	23	100_{\circ} 00

Early complications included gaping of stitch line in 6 cases (without exposure of bone). There was persistent discharge after operation in 4 cases while in 1 patient bone was exposed after the gaping of stitch line. One patient expired due to flaring up of septicemia and bacterimia post operatively. Sequestra formation and recurrence of infection was seen in one case. One patient had pathological fracture after 2 months of surgery (Table XVIII).

Table	XVIII	2	Showing	various	complications.
-------	-------	---	---------	---------	----------------

Com	plications No.	of	Cases
۸.	EARLY		
	Sloughing of skin with exposure of bone	1	
	Gaping of stitch line (without exposure of bone)	6	
	Persistant discharge after operation	4	}
	Neurovascular	ag	•
в.	And the state of t		
	Recurrence of infection	1	L
	Sequestrum formation	1	
	Pathological fracture	1	
		ninghating dividual for	

Note : One patient was expired due to post-operative flaring up on septicemia and bacterimia.

was the blockage of suction tube. This problem was maximum in first two days after the surgery. This problem was solved in later cases by keeping the irrigation flow fast for first 2 days and by reversing the irrigation flow. However, this problem was faced in as many as 5 cases. None of the case required discontinuation of irrigation suction technique. Leaking was seen in 4 cases. In one case we had to discontinue the suction irrigation due to gapping of stitch line and exposure of bone.

The problem of swelling of the limb due to collection of irrigation fluid was secondary to the plugging of suction tube. This problem was faced in 5 cases where there was blocking of suction tube.

In 1 case the tube was pulled out during ist post operative day and was reinserted after reopening the wound in operation theatre (Table XDXA).

Table XIXA: Untoward incidence of suctionirrigation technique.

Untoward incidence	Mo.of Cases	Irrigation discontinued
Leaking	4	1
Plugging of tube	5	49
Swelling	5	dep
Tube pulled out	1	disp

In 16 (69.56%) cases there was no significant problem during the period of irrigation and suction while in 7 (30.44%) cases one or more problems were faced shown in table XIXB.

Table XIXB :

Problem in irrigation suction technique	No.02 cases	Percentage
Present	7	30,44
Absent	16	69.56
Total	23	100.00

from 6-21 days with the average of 13.17 days. Irrigation technique was continued for 6-10 days in 6(26.09%) cases. In 11(47.81%) cases it was continued for 11-15 days and in 5(21.74%) cases for 16-20 days while in one case(4.36%) it was continued for more than 20 days(Table XX, Fig. 4).




FIG. 4. DURATION OF IRRIGATION.

Table XX : Duration of irrigation.

nurati	to ac.	irrigation	No. of	CASOS	Percentage
	479				
0 -	-		*		ajeto
6 ~	10			5	26.09
11 -	15		1:	L	47.81
16 -	20			5	21.74
7 2	0		3	1.	4,36
	Total	et efember i det mastelle ver tipelheide um eige vergleidigt velgtelle protes i med det element	2	iliyakida sengaya palaman da sapada Arrishida ili	100.00

irrigation ranged 8.4 lit to 27.3 lit with the average of 17.7 lit. In 2(8.70.) cases 5-10 lit normal saline was used and in 5(21.74%) 10-15 lit. saline was used in each case. In 7(30.42%) cases 15-20 lit. saline was used. 20-25 lit. saline was used in 3(34.78%) cases while 725 lit. saline was used in one patient(4.36%) (Table XXI).

Table XXI: Total amount of saline used for irrigation.

Amount (Lit.)	No.of cases	Percentage
5- 10	2	8.70
10 - 15	5	21.74
15 - 20	7	30.42
20 - 25	8	34.78
7 25	3.	4.36
recommendation of the second o	23	100.00

Amount of the normal saline used for irrigation in different cases ranged from 1120 ml to 1600 ml per day

with the average of 1367.8 ml per day. In 2(8.70%) cases 1100-1200 ml saline per day was used. 1200-1300 ml saline per day was used in 6(26.09%) cases. 1300-1400 ml saline per day was used in 10(43.48%) cases for irrigation while in 5 cases more than 1400 ml saline was used per day for the purpose of irrigation (Table XXII).

Table XXII: Average amount of saline used for irrigation per day.

Amount of saline (ml)	No. of	Percentage
1100 - 1200	2	8,70
1200 - 1300	6	26.09
1300 - 1400	10	43.48
7 1400	\$	21.73
zotal	entication - the second control of the control of t	100.90

Out of all 303 days, 257(84.82%) days combination of crystalline penicillin and Gentamicin was used followed by the combination of Streptomycin and cephasoline for 21(6.93%) days while combination of ampicilline and Gentamicin was used for 14(4.62%) days and for 11(3.63%) days the combination of Crystalline penicillin and chloromphenical was used.

Crystalline penicillin and Gentamicin combination was used in 21 cases followed by Crystalline penicillin and chloromphenical combination in 3 cases. In 2 cases the combination of ampicilline and Gentamicin was used

and the combination of Streptomycin and cephanoline was used in 1 case (Table XXIII, Fig. 5).

Table XXIII : Antibiotic used for irrigation.

Table XXIII Antioaseas	efficie effection can annue.	afficient em com Silla co.	
Antibiotics used	RG 6	No.of days	Percentage
Crystalline penicillin	21	257	84.82
Ampicillin + Gentamicin	*	14	4.62
Crystalline Penicillin + Chleremphenicol	3	11	3.63
Streptomycin + Caphasoline	1	21	6.63
		303	100.00

The change of the antibiotics was required for irrigation in 4 (17.39%) cases while in 19(82.61%) cases it was not required (Table XXIV).

Table XXIV : Change of antibiotic during irrigation.

	No.of cases	Percentage
Antibiotic changed	A service of the serv	17.39
Antibiotic not changed	19	82,61
TO C B L	23	100.00

Duration of follow up

ranged from 4 to 11.5 months with the average of 6.76 months. Twelve cases (52.17%) were followed up for the pariod of 3-6 months. For 6(26.09%) cases it was 6-9 months while for 5(21.74%) patients it was 9-12 months (Table XXV).




FIG: 5 ANTIBIOTICS USED FOR IRRIGATION.

Table XXV : Duration of follow up.

		tion ths)	No.ef Cases	Percentage
0	esto	3	- Million	
3	956	6	12	52.17
6	400	9	6	26.09
9	1986	12	5	21.74
an earl market	Tot	tal	23	100.00

The result was excellent in 14 (60.87%) cases while good in 5(21.74%) cases. However results were poor in 4 (17.39%) cases (Table XXVI).


Table XXVI : Showing results.

Result	No.of cases	Percentage
Excellent	14	60.87
Good	\$	21.74
Poor	4	17.39
Total	23	100.00

Pationt No. 1




1. Pre-op. A.P. and Lat. Skiagram of Lt arm showing sequestrum inside the cavity with subperiosteal new bone formation of Humerus.




 Post=Op A.P. and Lat. Skiagram of LA arm showing saucerized cavity in humerus with both irrigation and suction tubes in place.


3. Fellow-up A.P. and Lat. skiagram of Lt arm showing obliteration of cavity with marrow cavity patient on both sides (Table 3 months after operation).




Patient No. 2



 Pre-op, A.P. and Lat. skiagram of Rt forearm showing sequestrum inside the cavity with subperiosteal new bone formation in Radius.




 Post-op. A.P. and Lat. skiagram of Rt forearm showing saucerized cavity with both irrigation and suction tubes in position.



3. Follow up A.P. and Lat. skiagram of Rt forearm showing no evidence of residual infection with reduction in cavity size. (taken 4 months after operation).

Patient No. 3



1. Pre-op. A.P. and Lat. skiegram of Rt thigh showing sequestrum with old united pathological fracture of famus.




 Post-op. A.P. and Lat. skiagram of Rt thigh showing both irrigation and suction tubes in position with limb in POP cost.


3. Pollow-up A.P. and Lat. skiegram of Rt. thigh showing no evidence of residual infection (taken 6 months after operation).



Patient No. 4




1. Pre-op. A.P. and Let. skiagram or Rt leg showing thickening and sclarosis of lower \/3 of tibia with closes.



 Post-op. A.P. and Lat. skiagram of Rt leg showing saucerised cavity with both irrigation and suction tubes in place.

respectivity to



3. Follow-up A.P. and Lat. skiagram of Rt. leg showing reduction in cavity size in tibia and no evidence of residual infection (taken 5 months after operation).

DISCUSSION

Present study "To evaluate the effect of continuous irrigation and suction technique by using antibiotic solution in the treatment of chronic osteomy myelitis" was carried out on 24 cases of chronic osteomyelitis of different bones. Along with above technique conventional methods in the treatment of chronic osteomyelitis, like squeerimation, sequestrectomy were also used whenever needed.

Vesty and waldvogel (1980) observed that haematogenous esteomyelitis is most often seen in childhood. Bajaj et al (1981) found the incidence of haematogenous esteomyelitis 63.33 percent.

In the present study incidence of haemategenous esteomyelitis was 87.50 percent. This is because that most of the patients in our study belongs to lower age croups.

In the present series the patients are ranged from 7-50 years with a mean age of 19.79 years. About two thirds (67.67%) of the cases were below 20 years; being almost same as observed by Rao and Sahu (1978) where 66.67% patients were below 16 years. Hany workers observed that osteomyelitis is common in younger age group (Vesty and Waldvegel, 1980). In the series of Bajaj et al (1981) 53.3% patients were in the age group of second decade. This higher incidence of osteomyelitis in younger age group can be ascertained because of haemategenous

mode of spread which is also more common in present series. The another explanation can be the pattern of circulation in the metaphyseal region and greater liability to trauma. (Mercer, 1959).

Out of 24 cases, 21 (87.5%) cases were males and 3 (12.5%) cases were females, this is due to greater susceptibility of males to injury (Mercer; 1959). Similar observations were found in many other series. Males were 77% in the series of Rao and Sahu (1978), 73.3% in the series of Bajaj et al (1981) and 70.36% in the series of Clauson et al (1973).

Out of 24 cases, 18(75%) cases belonged to lower or lower middle socio-economic class. This is because of poor socio-economic conditions of most of population living in Bundelkhand region.

particularly lower extremities (Waldvogel et al, 1970 and Edeiket et al, 1973). In the present series benes of lower extremities were involved in 75% cases with commonest being femur (45,83%) followed by tibia(25%).

Dajaj et al (1981) reported incidence of osteomyelitis in femar and tibia as 37.9% and 27.6% respectively. Similarly many other workers reported that femar and tibia were most common bones involved in osteomyelitis (Taylor and Handsley, 1970; Letts and Mong, 1975; Classon et al, 1973; and Michelinakis, 1972).

In our study duration of illness ranged from 1.5 months to 8 years with the average of 17.88 months. In the series of classon et al (1973) average duration of illness was 2.2 years. However in the study carried by Davis et al (1986) average duration of illness was 8.9 years. This would have been because majority of the patients in present study were of younger age group.

Turek (1984) mentioned sinuses as very common clinical feature in chronic osteomyelitis. All cases except one (95.83%) in present series had sinuses.

by gram positive organism, most common being staphylococcus aureus (Aegerter et al, 1975; Waldvogel and Vasty,
1980). Relly (1977) reported that predominant organism
cultured was staphylococcus sureus in about 40% patients
of chronic osteomyelitis. In our study staphylococcus
aureus was the causative organism in 41.65% cases. In
the series of Davis et al (1986) staphylococcus aureus
was isolated in 44.74% cases.

in 2 (8.34%) cases. Kelly (1977) reported incidence of mixed infection 18%, Gram negative becilli were present in 8(33.33%) cases. This is in line with observation of Kelly et al (1973) that there is a trend of increasing incidence of gram negative bacilli cultured from saucerized osteomyelitis cavities. Identical findings were reported by Bajaj et al (1981). Out of 18 cases of

chronic osteomyelitis gram negative bacilli found in 3(16.7%) (Rac and Sehu, 1978). Gram negative organisms were isolated either alone or in combination in as many as 40% patients (Clawson et al. 1973).

The our series the pus culture from the wound was sterile in 6 cases. Most of the patients had under gone some sort of surgical procedure and almost all of them were receiving some antibiotic before they were subjected to suction-irrigation. This might be the probable cause of absence of organism in culture (Bajaj et al, 1981).

contamination of suction irrigation system by superinfection is one of the most common complication which occur either on opening the system with use of syringe, change of bottles, or bags of irrigant solution (Clawson et al, 1973). Superinfection was detected by culture of drainage fluid. When organisms cultured are other than the preoperative organisms than it means that superinfection has taken place.

infection. This is identical (33.3%) to the series of Letts and Wong (1975). The organism most commonly isolated was pseudomonas (16.67%). In the study of Letts and Wong (1975) Pseudomonas was the commonest organism causing superinfection. This is because the gram magative organisms are commonly responsible for hospital acquired infection (Griego, 1972).

Most frequent problem seen was the blockage of suction tube in 5(21.74%) cases. Constant attention was demanded for first 2 days because during this period problem of elogging was maximum. To avoid this problem we kept the flow rate high for the first 2 days and reversed the suction irrigation system done periodically every few hours. Clogging of the tubes was common even when they were properly utilized, this was due to the fact that the pressure gradient decreases down the tube with the greatest flow occurring through the more proximal holes in the afferent tubes and the distal holes in the efferent tubes. Reversing the flow may not unclog all the holes. When clogging does not disappear the safest way to cleam the tubes by the use of a stripping technique. Lubricant was applied to avoid finger burns and to provide a fast high pressure bolus of solution down to the suction tube. During this manoeuvre the patients were explained that they may feel momentary discomfort as the fluid produces internal pressure effect (Clawson et al, 1973).

If the solution runs dry, the tubes could have blocked within minutes and frequently would not work even with stripping. It is never advocated to flush the system by syringing because ; even with the best aseptic precautions secondary contamination can occur. Clawson et al (1973) modified the system with a syringe mounted in place. However, we flushed the system with a syringe under full asaptic precautions, whenever stripping technique failed.

Problem of the clogging of the tubes was faced by many other workers (Letts and Hong, 1975; Dilmaghani et al. 1969; Smith-Peterson et al. 1945; Taylor and Maudsley, 1970). All the workers solved the problem by reversing the flow and by flushing.

The causes of the plugging of the suction tubes could be :

- 1. Too much "Suction pull" This tends to suck normal tissue down into the tube and thus seal it off.
- Excessive wound waste in the form of blood and necrotic material.
- 3. Too little suction. This allows necrotic material to collect in the tube leading to its block.

Too little suction may occur if the tube is displaced and perforated end comes too near to the skin exit, thus allowing an air leak. This can be prevented initially by having a long tract for the tube to traverse, to ensure that all perforations were within the tissues (Taylor and Maudsley, 1970).

Other problems were local pain and swelling. These were secondary to the plugging of section tube. As outflow is blocked and irrigation fluid continuously entering the cavity thus increasing the pressure inside and resulting local pain and swelling. Problem was automatically solved with the re-establishment of irrigation-suction system. Similar problems were seen by many other workers (Dilmaghani et al, 1969; Smith-

Peterson et al. 1945; Letts and Hong, 1975; Clawson et al. 1973; Taylor and Maudsley, 1970).

In the present series 5(21.74%) cases developed the problem of blockage of suction tube.

Leaking was another common problem in 4(17.36%) cases which can be explained by :

- Repeated clogging of the suction tube resulting to increased pressure inside the cavity and thus gapping of the stitch line.
- Displacement of the tubes thus bringing perforated ends too near to the skin exit, which could be avoided by tight anchorage and deeper insertion of tubes.

Many other workers faced the same problem and were able to solve in the same way (Amith-Peterson at al. 1945; Dilmaghani et al. 1949; Taylor and Mandeley. 1970; Clawson et al. 1973; Letts and Wong; 1975). In 1 case leaking was very much because of gapping of theskin and exposure of the bone and we had to discontinue the irrigation-suction technique.

In 1(4,36%) case patient himself accidently pulled the tube which was reinserted after re-opening the wound in operation theatre. This has also been reported by (Letts and Wong, 1975 and Michelinakis, 1972).

In the present study complications were seen in 7(30.44%) cases and remaining 16 (69.86%) cases were problem free.

In the present study duration of irrigationsuction technique ranged from 6-21 days with the average of 13.17 days. In about half of the cases (47.81%) irrigation was continued for 11-15 days. It was difficult to know how long the tubes should be left in the wound. We kept the tube inside till two consecutive cultures of drainage fluid found to be sterile. Clawson et al (1973) kept the tubes inside till the dead space obliterated. They took the help of sinography for determination of dead space. They thought that cultures of the egress fluid is not a satisfactory criteria for removal of the tubes as positive cultures may indicate cross contamination from technical breakdown in the system. Under these circumstances, the bacteria may be growing in the tube system and not be pathogenic in the host. In their series irrigation was continued for 5 to 40 days (80% of the tubes removed between tenth and twenthieth day).

Bajaj et al (1981) used same criteria for the removal of the tubes as used in our series and the duration of irrigation (4-14 days) was comparable to our study. Reo and Sahu (1978) also used same criteria for the removal for tubes.

duration of irrigation as 3 to 4 weeks. They explained this duration on the basis that at least this much time was required to organise the haematoma filled dead space with granulation tissue. However they have continued

irrigation for 1 week after 3 consecutive negative cultures have been obtained.

Letts and Wong (1975) used closed tube irrigation in their series for 4-40 days with average of 13.5 days. However in the series of Taylor and Maudsley (1970) irrigation was continued for upto 6 weeks.

In our study amount of normal saline used for irrigation in different cases ranged from 1120 ml to 1500 ml per day with the average of 1353.8 ml per day.
1300-1400 ml normal saline per day was used in 10(43.48%) cases. We kept the flow rate of the irrigation solution high. This not only helped to prevent sludging during the early part of the irrigation but also might be essential to the mechanism of function of the procedure (Dombrowski and Dunn, 1965). During the first few days of irrigation, a high flow rate was required to prevent clogging of the suction tube. This is because bleeding inside the cavity was significant and thus enhancing the chances of clogging of the tubes. This theory is supported by many other workers (Dombrowski and Dunn, 1965; Taylor and Maudsley, 1970 and Clawson et al. 1973).

Different workers kept the flow rate of the irrigation solution as follows :

- 1. Dombrowski and Dunn (1965) 2-3 Lt/day/tube
- 2. Compare (1962) 2 Lit/day
- 3. Michelinakis (1972) 2 Lit/day
- 4. Taylor and Maudsley (1970) 1000-1500 ml/day

5. Rao and Saha (1978)

1000-1500 ml/day

6. Bajaj et al (1981)

1000-1500 ml/day

7. Clawson et al (1973)

1-2 Lit/hr for first two days.

Amount of irrigation solution used by most of the workers ranged from 1-2 Lit/day. Only Clawson et al (1973) used the irrigation solution in very high amount but they did not used antibiotics in the irrigation fluid.

Antibiotics were used in the irrigation fluid according to the culture report of the drainage fluid.

Combination of crystalline penicillin and Gentamicin were used. Other antibiotics used were - Streptomycin,

Cophasoline, Ampicilline and Chloramphenicol. Basis

behind the use of antibiotics in the irrigation fluid was the delivery of Deffective antibiotics to the infected region in high concentration with minimum systemic distribution (Dilmaghani et al. 1969).

Besides the local antibiotics we also used systemic antibiotics.

Combination of local and systemic antibiotics was used by many other workers viz. Rao and Sahu (1978), Bajaj et al. (1981), Dilmaghani et al (1969), Compare (1962) and Smith-Petersen et al (1945).

However, many workers used detergent or wetting agent along with irrigation fluid (Compere, 1962; Michelinakis, 1972; Dilmaghani et al, 1969; Mitra and Grace, 1956-57).

our study ranged from 14-40 days with the average of 22.26 days. Fourteen (60.81%) cases had post-operative stay of less then 3 weeks. It indicates that post-operative stay with this technique was very brief. Same conclusion was drawn by many other workers (Smith-Petersen et al. 1945; Dombrowski and Dunn, 1965; Rao and Sahu, 1978 and Bajaj et al. 1981). However, Letts and Wong (1975) reported in their study that children treated with closed tube irrigation required a period of post-operative hospitalization almost twice as long as that of the control group. But difference in this study was that they used the technique for the treatment of acute esteomyelitis in children. Moreover they themselves emphasized that closed-tube irrigation is an excellent method of treating chronic osteomyelitis.

arief post-operative stay ensures excellence of the technique for primary healing of the wound.

In our study duration of follow up ranged from 4 to 11.5 months with the average of 6.76 months. Rao and Sahe (1978) followed the patient for almost same duration (7.9 months). Taylor and Maudslay (1970) followed their patients for at least 1 year. Dombrowski and Dunn (1965) followed their patients for the pariod of at least 6 months. Dilmeghani et al (1969) followed their patients for the pariod of 2 years.

Different workers used various criteria for the evaluation of the results. We used definite clinical and

radiological criteria which were same as used by Bajaj et al (1981) for the assessment of the results in three grades i.e. excellent, good and poor.

Excellent

When there was healing by primary intention without any sinus, infection, pain or local tenderness.

good

When there was delayed wound healing, occasional pain and tenderness but without discharge, sinus or radiological evidence of infection.

Poor

When there was persistant pain or discharge with radiological evidence of infection or formation of sequestrum.

With above criteria results in our study were as follows :

Excellent in 14 (60.87%) cases.

Good in 5 (21,74%) cases.

Peor in 4 (17,39%) cases.

Different criteria were used by other workers e.g. Rao and Sahu (1973) evaluated their results as "Successful" and "PAilure".

Mamy workers evaluated their results according to recurrence of infection. If no recurrence them they considered result as good (Compere, 1962; Michelinakis, 1973; Dilmaghani et al, 1969; Smith-Petersen et al, 1945; Letts and Wong, 1975; Taylor and Maudaley, 1970).

Demborwski et al (1969) encouraged with the short term results of closed irrigation-suction technique. They compare their short term results with these obtained by the open packing method in another series. Passage of 6 months without recurrence of drainage was taken arbitratily as the criterion for comparison.

Clawson et al (1973) in their series evaluated results by following 3 parameters : drainage, pain and toxic condition. The patient's condition on follow up was compared to their pre-operative condition and change noted in the 3 parameters and results were evaluated.

COMPARISON OF SUCCESS RATE

Assence	Year	TotalNo. of cases	Result (Success) in percent
Grace and Bryson	(1945)	3	100,00
Mitra and Grace (19:	57) (Calcutta)	95	64.21
	(Brooklyn)	45	69.00
Goldman et al	(1960)	5	100.00
Dombrowski and Dunn	(1965)	20	77.00
Dilmaghani et al	(1969)	24	87.50
Taylor and Mandsley	(1970)	12	66.70
Michelinakis	(1972)	12	83.33
Clawson et al	(1973)	176	74.00
Reo and Sahu	(1978)	18	77.70
Sajaj et al	(1981)	30	73.30
Present series	(1987-88)	23	82.61

The treatment of chronic osteomyalitis is vary cumbersome, time consuming and unsatisfactory till date which signifies from the fact that several attempts to cure it have been made starting from boiling water to various reconstructive operative procedures, but none is foolproof.

All conventional methods available at present had one common aspect i.e. removal of deed tissue and sequestra excision of sinuses and saucerisation. The dead space thus left behind is filled up by the haematoma which serve as potential site of the infection.

By using continuous irrigation-suction technique this problem is solved.

Moreover, effective entibiotics can be delivered to infected region in high concentration with minimal systemic distribution by adding them in irrigation fluid. Otherwise antibiotics could not reach at the site of infection because of the relative avascularity of bone and inability of antibiotics to transgress the area occupied by necrosed tissue and pus.

Suction technique includes - Almost complete primary closure of the wound which is not possible with other methods and thus resulting to a minimum loss of serum from the epithe-lial surface. This is important because the loss of serum protein can be considerable, as was demonstrated by Smith-Petersen et al (1945) by treating 1 case by the method of "Packing and wound open". The determined loss of serum

protein for a period of 24 hours amount for 36.9 grame; the average daily loss over a period of a week was 11.9 gms.

The infrequency of dressings appeals the patients and to the surgeon. Septic dressings are painful no matter how gently they are done, and they are bound to be time consuming.

The time for the complete healing of the wound is shorter and sear remain linear and smooth, instead of broad and puckered (Smith-Petersen et al, 1948).

Dombrowski and Duna (1965) reported requirement of irrigation system as follows:

- Air tight and water tight wound closure This prevents skin maceration, reintroduction of new organisms.
- 2. Placement of irrigation tubes wherever necessary.
- 3. Keeping multiperforated ends of the tubes short
 (3 of 4 inch) because when fluid is in motion there is
 a constant fall in hydrostatic pressure along the
 perforations distally (Binder, 1949). There is also
 a constant fall in negative pressure distally along
 such perforated suction tubes. When tubes are long
 clots form distally, migrate proximally, ultimately
 plugging the system.
- 4. Keeping the flow rate of the irrigation solution high this prevents plugging of tubes.
- 5. Adequate duration (2-3 weeks) of irrigation discontinued when 2 consecutive cultures become negative.

- 6. Use of appropriate systemic and local antibiotics according to culture reports.
- 7. Good nursings and house staff are essential because constant attention is necessary for the first 2 days and it is impossible for a surgeon to manage the suction irrigation system himself (Classon et al, 1973).

Smith-Petersen et al (1945) reported that :

- This is not a perfect method with 100% cures. There had been cases with local recurrence of sepsis, but these had been relatively few.
- The suction and irrigation system is not immune to clogging. But a change in the direction of the irrigation fluid and flushing the wound usually aliminate obstructions without much difficulty.
- Leaking is also a problem but this can be solved by air tight and water tight wound closure.

Closure of the wounds with irrigation and drainage were first reported by Smith-Petersen et al (1945). In 1956 Mitra and Grace treated esteomyelitis by instillation of penicillin and a detergent through catheter into the wound. In 1060, Goldman et al and in 1961 Mc Elvenmy described the use of a circulation suction method of treating infected wound and esteomyelitis. To improve the function of the system and to enhance the antibacterial effects of the solution, Compere (1962) recommended the addition of tylomapol (Aleyaire) to irrigating fluid. In

1965 Dembrowski and Dunn used streptokinase - Streptodornase (Varidase) and in 1969 Diffehani et al used chlorpactin XCB in the suction irrigation. Clawson et al (1965, 1973) modified the previous tubing system in closed suction irrigation technique.

Dombrowski and Dunn (1965) postulated the following explanation of why prolonged irrigation may seen to substitube atlasst in part even for inadequate debridement. Irrigation solution composed of Physiologic saline and antibiotics (Camotic pressure of about 30 mm of Hg) are hypertonic with respect to interstitial fluid (Comotic pressure of about 15 mm of Hg). There is an esmotic gradient of 15 mm of Hg. in: ducing a flow of fluid from tissue to wound eavity. The second factor which helps the flow of fluid is the temperature. The lower the temperature the greater the comotic pressure. The tissue temperature is about 37°C and the temperature of cavity is about 25°C (Temperature of irrigating system at room condition). Thus there is again a gradient favouring flow of fluid from local tissue to wound cavity. This flow brings fresh lymph and leucocytes into the area which have antibacterial preparty and help in obliteration of dead space.

Closed suction irrigation technique incorporates all previous principles of treatment, with the patient at bed rest and host defences mobilised. Radical debridement eliminates involved dead and devitalized tissue. Dead space heemstoma is controlled by irrigating solution by

Certain possible physiologic meghanism, as postulated by Dombrowski and Dumn (1965). Primary closure of skin and soft tissues prevent cross contamination and recontamination. The presence of antibiotics through irrigation solution aids local host mechanism. High concentration of local antibiotic infusion controls any septic process persisting in the area.

Poor result from the use of closed irrigation suction is usually from one or more causes as stated by Clawson et al (1965, 1973).

Inability to remove all the dead bone and surrounding scar tissue :

- Feilure to understand the principles of suction irrigation system.
- The advantages of suction-irrigation technique can be summarised as follows:
- Almost complete prediction that the operative wound will heal by primary intention.
- 2. Infrequency of dressing.
- 3. Reduction in the time for complete healing.
- 4. Minimum loss of serum.
- 5. No soiling of bed linen.
- 6. Comfort and lack of systemic reaction.
- Reduction in cross infection, hospitalization and post-operative isolation.

n. Metalic intramedullary fixation may be left in place or inserted at the time of operation.

Most important limitation with this technique is that it cannot be used in the treatment of osteomyelitis of short bones.

Though observation period is short and the material is small, but on the basis of results achieved and comparing the results with other similar series it is justified to recommend, the use of continuous irrejection suction by using antibiotic solution in the treatment of chronic osteomyelitis.

CONCLUSION

continuous irrigation-suction technique by using antibiotic solution is an excellent mode of treatment for chronic osteomyelitis, when used as an adjuvant to radical excision of unhealthy tissue and administration of effective antibiotics. Irrigation with antibiotic solution serve dual purpose. Firstly, it prevents heematoma formation and secondary high concentration of antibiotics reaches at local site.

In the present study following conclusions are drawn a

- Younger ones (\(\sigma 20 \) years) are affected by osteomyelitis more than older ones.
- Males are more commonly affected than females.
 Male, female ratio being 7:1.
- 3. Haematogenous spread is common mode of infection.
- 4. The prevalence of disease is commoner in lower socio-economic group as 75% cases are from lower and lower middle class.
- Lewer limb is more prone to disease rather than other sites.
- The usual clinical feature presented are bone pains, tenderness, pas discharge and non healing wounds.
- Radiological signs commonly seen are localised sclerosis, disfigured out line, destruction, periosteal reaction, sequestration and sinuses.

- 8. Staphylococcus aureus is commonest organism causing esteomyelitis.
- 9. Pseudomonas is commonly seen causing superinfection.
- 10. Common problems found in the present technique are leakage, plugging and swelling secondary to plugging. These problem can be easily controlled.
- 11. High rate of irrigation flow is required to prevent plugging. Air-tight closure of wound is necessry to prevent leaking.
- 12. 1100-1600 mo normal saline is the usual volume required for irrigation.
- Appropriate duration of irrigation should be
 2-3 weeks.
- 14. Crystalline penicillin and gentamicin are found to be most effective antibiotics in the treatment of osteomyelitis.
- 15. Redical excision of all devitalised soft tissue and dead bone followed by instillation of continuous irrigation and suction technique by using antibiotic solution has shown good or excellent results in \$2.61% cases with failure in 17.39% cases.
- 16. Incomplete removal of sequestrum, dead tissue, presence of superinfection, resistant infection are commonest causative factors for failure of technique.

BIBLIOGRAPHY

- i. Aegerter, E. and Kirkpatrik, J. (1975) : Orthopadic disease, 4th ed. Philadelphia, WB Saunders.
- Albee, P.H. (1933) : The treatment of esteomyelitis
 by bacteriophages, J.B.J.S., 15 : 58.
- 3. Andrew, J. Weiland; Pussel, J.; Moore and Rellin, K.;
 Demiel (1984): The efficacy of free tissue transfer
 in the treamtment of osteomyelitis. J.B.J.S., 66-A;
 No. 2, 181-193;
- 4. Ash, J.M. and Gilday, D.L. (1980): The futility of bone scanning in neonatal esteemyelitis. Concise communication. J. Nucl. Med., 21 : 417.
- 5. Seeker, R.O. and Spedare, J.A. (1978) : Treatment of orthopsedic infections with electrically generated silver ions. J.B.J.S., 60-A : 871.
- Baer, W.S. ((1931): Treatment of chronic osteomyelitis with magget, J.B.J.S., 13: 438.
- 7. Bajaj, A.S., Maini, P.S., Arora, D.R., Madgil, K.C. and Arora, B. (1981): Treatment of chronic osteomye-litis with continuous irrigation suction technique: Indian Journal Orthop, Vol. 15, No.2:211-2164
- 8. Berkowitz, I.D. and Wensel, W. (1980) : "Mormal" technetium bone scans in patients with acute osteo-myelitis. Am. J. Dis. Child, 132 : 828.

- 9. Bickel, N.H.; Bateman, J.G. and Johnson, W.R. (1953):
 Treatment of chronic basmatogenous esteomyelitis by
 means of saucerization and bone grafting surgery.

 Gynascol & Chatet, 96: 265.
- 10. Binder, R.C. (1949) : Fluid mechanics, ed. 2, New York, Prentice-Hall.
- 11. Bordon, L. and Buncke, H.J. (1983): Limb salvage achieved with tissue grafting. Orthop News, 5.
- 12. Bosch, P. (1981): Die Fibrinospongiosaplastic:
 Experimentalle untersuchungen und klinische Erfahrung
 Wiem klin Wochenschr (Suppl), 93(124): 1-26.
 (Eng Abstr.).
- 13. Bosch, P.; Braun, F.; Eschberger, J.; Holle J.J. (1976):

 Experimentalle untersuchungen über die Ammendungdes

 "Fibrinklebesystems" bei der knochenheilung G. Forum
 für experimentalle Medizin Osterreichischer

 Arstekongress Von Swieten Tegung.
- 14. Brantigan, O.C. and Owen J.C.(1942) : sodium tetradecyl sulphate used in the treatment of acute pyogenic emphyema of the pleural cawity. Bull. School Med. Univ. Maryland, 26 : 247.
- 15. Bryson, A.F.; Mandell, B.B. (1964): Primary closure after operative treatment of grees chronic esteemyemlitis. Langet, Vol. 1 : 1179.
- 16. Burri, C.; Passier, H.H. and Henkeneyer, H. (1973):
 Treatment of post-traumatic osteomyelitis with bone,
 soft tissue and skin defects, J. Trauma, 13: 799.

- 17. Burwell, R.G. (1964) : Studies in the transplantation of bone VII. The fresh composite homograft-autograft of cancellous bone : an analysis of factors leading to osteogenesis in marrow transplants and in marrow contemining bone grafts. J.B.J.S.(Br.) 46-B : 110-40.
- 18. Capitanio, M.A. and Kirkpetrik, J.A. (1970) 4/Rosntgen observations acute osteomyelitis. Am.J. Rosntgenol. Radium Ther. Nucl. Med., 108 : 488.
- 19. Clawson, B.K.; Davis, F.J. and Hansen, S.T. (1973) :
 Treatment of chronic osteomyelitis with emphasis on
 closed suction-irrigation technique. Clin. Orthop.
 and related research. No. 96 : 88-97.
- 20. Clawson, D.K. and Stevenson, J.K. (1965): Treatment of chronic osteomyelitis. Surgery, Gynse. & Chst. 120: 59.
- 21. Colemen , H.N.; Batman, J.C.; Dale, G.M. and Storr, D.E. (1946): Campellous bone grafts for infected bone defects: A single stage procedure. Surg. Gynaecol Chstet., 83: 392.
- 22. Compare, E.C. (1962): Treatment of esteomyelitis and infected wounds by closed irrigation with detergent antibiotic solution. Acta. Orthop Scand., 32 : 324.
- 23. Compere, E.C., Metzger, W.I. and Mitra, R.H. (1967):
 Treatment of pyogenic bone and joint infection by
 closed irrigation (Circulation) with non toxic
 detergent and one or more antibiotics. J.B.J.S.(Am.)
 49:614.

- 24. Dalinka, M.K.; Dennengerg, S.; Greendyke, W.H. et al (1971): Roentgenographic features of osseous coecidioidomycosis and differential diagnosis. J.B.J.S.; 53-A; 1187.
- 25. Dalinka, M.K.; Lally, J.F.; Koniver, G. et al(1975); The radiology of osseous and articular infection. CRC Crit. Rev. Clin. Radiol. Nucl. Med. 7:1.
- 26. Davis, J.C. (In press): Adjunctive hyperbasic owygen in chronic refractory esteomyelitis. Long term follow-up results (Letter) Clin. Orthop.
- 27. Davis, J.C.; Heekman, J.D.; Delee, J.C.; Buckwold, F.J. (1986) : Chronic non-haematogenous estecmyelitis treated with adjuvant hyperbaric enygen. J.B.J.s., 68-A, No. 8: 1210-1217.
- 28. De Oliveira, J.C. (1971): Bone grafts and chronic osteomyelitis. J.B.J.S.(Br.), 36: 584.
- 29. Dilmaghami, A., Close, J.R., Rhimelander, F.W. (1969):
 A method of closed irrigation and suction therapy
 in deep wound infections. J.B.J.S., Vol. 51-A, No. 2:
 323-342.
- 30. Dombrowski, B.T. and Dumn, R.W. (1965): Treatment of osteomyelitis by debridement and closed wound irrigation-suction. Cli. Orthop., 43 : 215.
- 31. Duman and Carrel (1917) : Cited by Rao and Sahu, 1978.
- 32. Edeiken, J. and Hodes, P.J. (1973) : Roembgen diagnomis of disease of bone. 2nd ed. Baltimore, Williams and Wilkens Company.

- 33. Engh, C.A.; Hughes, J.L. and Abrams, R.C. et al (1971): Osteomyelitis in the patient with sickel cell disease, J.B.J.S., 53-A : 1.
- 34. Friedman, S.A. and Rakow, R.B. (1971) : Osseous lesion of the foot in diabetic neuropathy. Diabetes, 20:302.
- 35. Fitsgerald, R.H.Jr.(1979): The epidemilegy of noseconial infections of the misculeskeletal system. In hospital associated infections in the general hospital population and specific measures of control. Edited by Dieter Groschel, Vol. 3, 25-35, New York, Dekker.
- 36. Ger, R. (1970): The management of open fractures of the tibia with skin loss. J. Trauma, 10: 112.
- 37. Ger, R. (1972) : Surgical management of ulcerative lesions of the leg. Curr. Probl. Surg., 1.
- 38. Ger. R. (1977): The closure of defects in the lower extremity by muscle flaps. In convers, J.M. (ed.)
 Reconstructive plastic surgery Philadelphia.
 Saunders, 3549.
- 39. Goldman, M.A.; Johnson, R.K. and Grossberg, N.M.

 (1960): A new approach to chronic esteomyelitis.

 Orthopaecies.
- 40. Gerdon, L. and Buncke, H.J. (1983) : Limb salvage achieved with tissue grafting. Orthop. News, 5.
- 41. Grace, EJ. (1955): Topical streptchydrasid treatment of a tuberculus sinus. Antibictic medicine 1: No. 2, 104-105.

- 42. Grace, E.J. and Bryson, V. (1945): Topical use of concentrated penicillin in surface active solution. Achives of surgery, Vol. 50: 219-222.
- 43. Grace, E.J. and Sryson, V. (1947) : Chronic osteomogelitis in war wounds : A report of two veterans discharged with intractable osteomyelitis and successfully treated with local penicillin-detergent therapy.

 M.Y. State J. of Hed. 47, No. 27 : 2204-2205.
- 44. Grace, E.J. and Bryson, V. (1950): Modern treatment of chronic osteomyelitis with topical detergent antibiotics therapy. Surg. Gynas. & Obstet. Mcl. 333-342.
- 45. Grieco, M.H. (1972): Pseudomonas arthritis and osteomyelitis. J.B.J.S., 54-A: 1693.
- 46. Owinn, J.J.; Free, F.A. et al (1972) : Rediologic case of mometh. Am. J. Dis. Child, 124 : 91.
- 47. Hagen, R. (1978) : Osteomyelitis after operative fracture treatment. Acts. Orthop. Scend. 49: 542.
- 49. Hall, J.E. and Grossfield, L. (1966): The management of acute hematogenous osteomyelitis. Applied Thorap. 8: 331-335.
- 49. Hirschmann, J.V. and Everett, E.D. (1976) : Candida vertebral osteomyelitis case report and review of literature. J.B.J.S., 58-A: 573.
- 50. Howie, D.W., Savage, J.P., Wilson, T.G., Peterson, D.W. (1983): The technetium phosphate bone scen in the diagnosis of osteomyelitis in childhood.

 J.B.J.S., 65-A(4): 431-437.

- 51. Hazlette, J.W. (1954): The use of cancellous bone grafts in the treatment of subscute, chronic esteomyelitis. J.B.J.S.(Br.), 35: 584.
 - 52. Kelly, P.S. (1927): Infections of bones and joints in adults patients. Instructional course lectures.

 American Academy of orthopaedic Surgeons, 26:3-13.
- 53. Kelly, P.J.; Wilkoske, C.J.; John, A. (1973):

 Comparison of gram negative bacillary and Staphyloeccal osteomyelitis of femur and tibia. Clinical
 Orthopaedics and related research, 96 : 70.
- 54. Kido, D.; Bryan, D. and Halpern, M. (1973) : Haematogenous esteemyelitis in drug addicts. Am. J.
 Roentgenol, 118: 356.
- 55. Komlewski, K., Hamel, J., Harbison, S. (1983) : J. Yu. Pediatr. Radiol., 13 : 130-136.
- 56. Kuha, J.P. and Burger, P.E. (1979) & Computed tomographic diagnosis of osteomyelitis. Radiology, 130:503.
- 57. Lack, W.; Bosch, P.; Arbes, H. (1987) : Chronic osteomyelitis treated by cancellous homografts and fibrin adhesion, J.B.J.S., 69-B, No. 2 : 335-337.
- 58. Lawer, R.B. and Eyring, E.J. (1972) : Intermittent closed suction-irrigation treatment of the osteomye-litis. Clin. Ortho. 88 : 80.
- 59. Letts, R.M. and Wong, E. (1975): Treatment of acute osteomyelitis in children by closed tube irrigation:
 A reassessment: The canadian Journal of Surgery.
 Vol. 18: 60-63.

- 50. Lieutement Colonel Marvin P. Knight and Major George O Wood (1945) : Surgical obliteration of bone cavities following traumatic osteomyalitis.
- 51. Mathes, S.J.; Alpert, B.S. and Chang, Ning (1982) :
 Use of the muscle flap in chronic esteomyelitis,
 experimental and clinical correlation. Plast and
 reconstr. Surg., 69 : 815-828.
- 62. May, J.W.Jr.; Gallico, G.G. III and Lukesh, P.N.

 (1982) : Microvascular transfer of free tissue for
 closure of bone wounds of the distal lower extremity.

 New England J. Med., 305 : 283-257.
- 63. May, J.W.Jr.; Savage, R.C. (1983): Microvascular transfer of latissimus muscle with skin graft for correction of chromic bony wounds. Read at the annual meeting of the International Society for Reconstructive Microsurgery, New York, N.Y.
- 64. Mc Elvenny, R.T. (1961): The use of closed circulation and suction in the treatment of chronically infected, acutely infected and potentially infected wounds. The Am. J. Orthopedics, 86-87 and 154-160.
- 65. Mercers W. (1959) : Orthopsedic Surgery, V ed., Edward Armold (Publishers) Ltd., London.
- 66. Michelinakis, E. (1972) : Treatment of chronic osteomyelitis with the continuous irrigation-suction method. Acta Orthop. Scandinav., 43 : 25-31.
- 67. Mitra, R.N. and Grace, E.J.: Further studies on the treatment of chronic osteomyelitis with topical detergent-antibiotic therapy. Antibiotic annual,

- 1956-57 Medical Encyclopedia. Inc. New York, N.Y. 455-466.
- 68. Morrey, B.P.; Dunn, J.M.; Heimbach, R.D. and Davis, J. (1979) : Hyperbaric oxygen and chronic osteomyelitis. Clin. Orthop, 144: 121-127.
- G9. Morrey, B.F., Fitzgerald, R.H. Jr., Kelly, P.J., Dobyns, J. II, Washington, J.A. II (1977) a Diphtheroid osteomyelitis. J.B.J.S. (Am.) Vol. 59 : \$27-530.
- 70. Mund, D.J. (1980): Pyogenic vertebral esteomyelitis: Manifestation of bacterial endocarditis, N.Y. State
 J. Med., 80: 980.
- 71. Marry, R.O. and Jacobsen, M.C. (1977): The radiology of skeletal diagnosis. 2nd ed. New York, Churchill Livingstone.
- 72. O'Riordan; Colm; Adler, J.L.; Banks, H.H.; and Finland Maxwell (1972): Wound infections on an erthopaedic service a prospective study. Am. J. Epidemiol, 95: 442-450.
- 73. Orr, H.W. (1927): Treatment of osteomyelitis and other infected wounds by drainage and rest. Surg. Gynae and Obst., 45 : 446-464.
- 74. Papimeau, L.J. (1973) : L' Excision-grafte avec fermeture retord'ee deliberee dans l'esteomyelite chronique Nouelle Presse Medicale, 2 : 2753-2755.
- 75. Polo, G.V. (1968) : Compare's method modified for treatment of tuberculosis of bone and joints. In

- proceedings of the association of bone and joint surgeons. Clin. Orthop. 58: 296.
- 76. Prigge, E.K. (1946): The treatment of chronic osteomyelitis by the use of muscle transplant or iliac graft. J.B.J.S., 28: 526.
- 77. Rae, P.T. and Sahu, P. (1978): Treatment of chronic ostcomyelitis with closed irrigation and suction technique: Indian Journal of Orthop. Vol 12 No. 2: 180-184.
- 78. Reynold, J. (1977): Radiologic manifestations of sickle cell haemoglobimopathy. J.A.M.A., 238: 247.
- 79. Rhinelander, F.W. (1975): Minimal internal fimation of tibial fractures: Clinical Orthopaedics and related research, 107: 188-220.
- 80. Riegler, H.F.; Goldstein, L.A. and Betts, R.F. (1974):
 Blastomycosis esteomyelitis. Clin. Orthop. 100:225.
- 81. Rewling, D.E. and Kirkealdy (1970): Further experience in management of chronic esteemyelitis.

 J.B.J.S., 52-B: 302-307.
- 82. Salahuddin, N.I.; Madhavan, J.; Fisher, E.J. et al (1973) : Pseudomonas esteomyelitis, Radiologic features. Radiology, 109 : 41.
- 83. Smith-Paterson, M.N., Larson, C.B. and Cochran, W.

 (1945): Local chemotherapy with primary closure of
 septic wounds by means of drainage of irrigation
 canulae. J.B.J.S., 27: 562.
- 84. Stark, W.J. (1946) : The use of pedicled muscle flaps in the surgical treatment of chronic osteomyelitis

- resulting from compound fractures. J.B.J.S., 28:343.
- 85. Stuart, A. Green and Margaret, J. Ripley (1984) : Chronic osteomyelitis in Pin tracks, J.B.J.S., 66-A: 1092-1098.
- 86. Svirsky-Fein, 5.; Langen, L.; Milbauer, B. et al (1980) : Neonatal Osteomyelitis caused by candida trepicalis. J.B.J.S.; 61-A : 455.
- 87. Taylor, A.R.; Maudeley, R.H. (1970) : Instillationsuction technique in chronic esteomyelitis. J.B.J.S.
 Vol. 52-B. No. 1 : 88-92.
- 88. Turck, S.L. (1984) : Orthopaedics : Principles and their application IV Ed. Vol. 1, J.B. Lippincott, Company, Philadelphia, : 262.
- 89. Ude, B. Heyme, Hisharm, K. Taminir David A Eschenbach
 Paul G Ramsey, David C Figge (1984) : Osteomyelitis
 Pubis after radical Gynecologic operation :
 Obst. & Gyn. Supple. 63 : 478-538.
- 90. Vivian W. Wing; R. Brooke Jeffrey, Jr.; Michel P.; Freierle Clyde A Helms and Peter Trafton (1985) : Ch. Osteomyelitis examined by computed tomography Radiology, 154 : 171-174.
- 91. Wajih Al-Sheikh; George, N. Siakianakis; Mmaymmeh, W.; Mukil Hourani; Heal, A.; Dungan, R.C.; Burnett, A.; Fuad-S Askhan; Aldo, N. Scrofini (1985); Subscute and chronic bone infections; Diagnosis using In 111. Ga 67, Tc 99 MDP, bone scintigraphy and radiography. Radiology, 155: 501-506.

- 92. Wald, E.R.; Mirro, R. and Gartner, J.C. (1980) :
 Pitfalls in the diagnosis of acute osteomyelitis
 by bone sean. Clin. Pediatr., 19 : 597.
- 93. Weldwogel, P.A.; Medoff, C. and Swartz, M.N. (1970);
 Osteomyelitis a review of clinical features,
 therapeutic consideration and unusual aspects
 (First Part) N. England J. Med., 282 : 198.
- 94. Waldwogel, F.A. and Vasey, N. (1980) : Osteomyelitis : The past decade. N. Engl. J. Med., 303 : 360.
- 95. Webester, D.A. (1981) : Silver enode treatment of chronic osteomyelitis. Clin. Ortho. 161 : 105-114.
- 96. Wiessemann, G.J.; Wood, V.E.; Kroll, L.L. (1973):

 Pseudomonas vertebral osteomyelitis in herein addicts,

 reports of five cases. J.B.J.S.(Am.) Vol.55:1416-1424.
- 97. Willenegger, H. and Roth, W. (1962) : The antibacterial irrigation drainage of surgical infections.
 Deutsch, Med. Wechr., 87 : 1485.
- 98. Woolfenden, Pitt, M.J., Durie, B.G.M., Moone, T.C. (1980) : Comparison of bone scintigraphy and radiography in multiple myeloma. Radiology, 134:723-728.



APPENDIX

TO EVALUATE THE EFFECT OF CONTINUOUS IRRIGATION AND SUCTION TECHNIQUE BY USING ANTIBIOTIC SOLUTION IN THE TREATMENT OF CHRONIC OSTEONYELITIS

Case No.

MRD No.

Mame

Age/Sex

Ward/Bod

Father's Ausband's Name

Cooupation

Address

DOA

DOD

Diagnosis

Consultant

Chief Complaints

History of present illness

History of Past Illness

Pamily History

Personal History

Socio-economic status

Ceneral Examination

Appearance Anaemia

Duilt Cyanosis

Pulse Clubbing

Taundice Jaundice

R/R Ondema

Lymph nodes Ceneral .

Local

Systemic Examination :

CVS

CMS

Resp.

Abdomen

Local Examination

- 1. Number of healed sinuses.
- 2. Number of discharging sinuses.
- 3. Operation scar mark.
- 4. Movements

Joint Above Joint Below

Abduction

Adduction

Medical rotation

Lateral rotation

flagion

Extension

Circumduction

Inversion

Eversion

5. Nerves -

Radial N.

Median N. Ulnar N.

Sciatic N.

Lateral Popliteal N.

6. Vascular Supply -

Brachial artery

Radial artery

Popliteal artery

Dorselis pedis artery Posterior tibial artery

- 7. Bony Thickeaing
- 8. Tenderness

Radiological Examination

Periesteal reaction Sequestrum

Destruction Sclerosis

Mottling Thickening

Cavity

Investigations

Blood TLC DLC : P L E

M

ISSR HD

Urine Albumin sugar M/E

Blood Urea

Blood Sugar : P

PP

Pus Culture & Sensitivity

Treatment

- 1. Pre-operative -
 - · ATE
 - . Tet. Vac.
 - . Antibiotic- Total No. of days : (1)

(11)

2. Operative Treatment -

- . Date of Operation
- . Type of Operation

Surgeon

Assistants

Anaesthesia

Op. Steps

Remarks

3. Post-Operative Treatment-

. Date of removal of stitches (i)

(11)

- . Type of immobilisation (A) Postoperative
 - (B) At the time of Discharge.
- . Duration of immobilisation (A)

(B)

Remarks

900 min	10. 1		uste di	South resting
Po	4	LON		

(i) Clinical Examination Date:

(11) X-ray

(iii) Movement - Above Joint

Below Joint

*Ngalija i malija njalata			
(1)	Healing		
(2)	Pain		
(3)	Tendernoes		
(4)	Sinus		
	Residual Infection		

Result

SULTANIS /

TRRIGATION SUCTION DE TAYES

82. No.	Date	Pluid	Amount	Amtibiotic used	Systemic Antibiotic	Permarks

64						
643						
**						
w						
võ			·			
co°						
•						
10						
and and						
75						
617) prof						
*						
uî M						

m	N/A	20	(Prop.	00	Q1	0	कनी कनी	P9	dod and	44	the g
Miles and the supplemental states are supplemental states and the supplemental states and the supplemental states and the supplemental states are supplemental states and the supplemental states and the supplemental states are supplemental states are supplemental states and the supplemental states are suppleme	The same of the same of				STATE OF STREET, SHAPE				The state of the s	The same of the same of	

Organian

Centerycth

0

Stroptosycin

Yerranyctn


Children conventin

Erythromycln

Closectile

Capitol extra

Polymyxin-E



continuous irrigation-suction technique by using antibiotic solution in the treatment of chronic osteomyelitis
was carried out on 24 patients of proved chronic osteomyelitis of different bones, admitted in Orthogodic ward
of M.L.B. Medical College, Hospital, Jhansi between July,
1987 and June, 1988. Out of 24 patients 1 patient died
post-operatively due to flaring up of septicemia and
bectarimia.

The criteria for selection of cases were pain, discharge, tenderness and thickening with radiological evidence suggestive of chronic osteomyelitis.

examined clinically and radiologically. Various relevant investigations were done prior to surgical intervention. According to pus culture and sensitivity report proper antibiotic coverage was given and surgery was performed in the form of saucerization, sequestrectomy, curettage, excision of sinus tract followed by placement of irrigation and suction tubes inside the cavity under general or regional anaesthesia. Irrigation-suction technique was checked per-operatively.

The limbs were properly supported either in slab or splint and patients were shifted to post operative ward. Patients and irrigation suction technique were closely watched post-operatively and any problem if arised was solved accordingly.

After stitch removal pain and tenderness of were reduced considerably in most/the cases.

These patients were followed up at regular intervals to evaluate clinically and radiologically. On the basis of these observations and evaluation results were categorised in the following three grades i.e. excellent, good and poor.

Excellent - when there was healing by primary intension, no sinus or residual infection. No pain or local tenderness, no radiological evidence of infection.

Good - when there was delayed wound healing or occasional pain and tenderness but no discharge or sinus and no radiological evidence of infection.

Poor - when there was persistent pain or discharge or radiological evidence of infection or formation of sequestrum.

In present study of 24 cases of chronic coteomyelitis most of the patients were in the age group of 11-20 years (with mean age of 19.79 years). Marked male predominance (87.50%) was observed due to their more outdoor activities. Commonest mode of spread of osteomyelitis was haematogenous.

Bones of lower limbs involved more (75%) with femmer as commomest site (45.83%). The duration of illness in these patients ranged from 1/2 months to 8 years. It was observed that most of the patients had

undergone some sort of treatment before this study (chemotherapy alonge or alongwith incision and drainage).

Sclerosis, thickening, destruction, sequestrum, periosteal reaction and cavities are the common radiological features. Staphylogoccus aureus was the commonest organism isolated from 10(41.65%) cases followed by Pseudomones (12.50%). Commonest organism responsible for superinfection was Pseudomones (16.67%). 8(33.3%) cases showed superinfection while in 16(66.7%) cases there was no superinfection.

Along with irrigation and suction one or more of the following procedures was carried out vis. curettage, syringectomy drill holes, saucerization, sequestrectomy.

Gaping of the stitch line, sloughing of skin and persistant discharge after operation were the complications observed.

includes plugging of suction tubes, leaking, swelling and pain at local site. All the problems were controlable. In 7(30.44%) cases these problems were seen while rest 16 (69.56%) cases were problems free.

We kept irrigation flow rapid for first 2 days to avoid the problem of clogging, 1200 -1600 ml normal saline was used for irrigation per day. Duration of irrigation ranged from 6 to 21 days. Crystalline penicillin and gentamicin were the choice of entiblotics used in irrigation fluid.

ranged from 14-40 days with the average stay of 22,26 days. Fourteen (60,87%) cases remained in the hospital for less than 3 weeks.

In present series, follow up of patients was ranged from 4 to 11.5 months with the average of 6.76months.

In this study results were excellent in 14 (60.87%) cases, good in 5(21.74%) cases and poor in 4(17.39%) cases. Thus results were fevourable in 82.61% cases.

In one case irrigation was discontinued backwase of sloughing of the skin and exposure of the bone while in one case recurrence of infection was seen.

Causes of the poor results includes inability to excise all deed and devitalised tissue and technical failures.

on the basis of this study, it is concluded that continuous irrigation suction technique by using antibiotic solution is an excellent method to treat chronic osteomyelitis, when used as an adjuvant to radical excision of unhaelthy tissues and administration of effective antibiotics.

Advantages of ixrigation suction technique are:

- 1. Hasmatama formation in the dead space is prevented thus minimising the chances of reinfection.
- 2. High concentration of antibiotics reached at local
- 3. Primary closure of the wound is almost certain.
- 4. Minimal loss of the serum.
- 5. Post-operative hospital stay is of short duration.